INDEX TO VOLUME XXIII

New species in blackface type

ABBOTT, E. V., Physiologic forms of Colletotrichum falcatum Went., 557–559

Abies concolor, attacked by Rehmiellopsis bohemica, 108–109

Abnormalities, leaf, in tomato mosaic, 26

Abutilon, seed transmission of virus causing variegation of, 20

Acer, rubrum, Nectria canker of, 36; spicatum, Nectria canker of, 36

Acetic acid, treatment with, for control of gladiolus scab, 808

Acid, production by Phytophthora nicotianae, 610

Acids, effect of, on Corticium koleroga, 880

Acta Phaenologica, new journal of phenology, review of, 316–317

Aecia of spruce needle rusts in Minnesota, 613, 614

Air, upper, spores in, 23

ALBERT, W. B., see ARMSTRONG, G. M.

Alfalfa, potato-leaf-hopper injury of, 19

ALLEN, RUTH F., A cytological study of teliospores, promycelia, and sporidia in Puccinia malvacearum, 572–586; Heterothallism in Puccinia graminis, P. coronata, and Melampsora lini, 4; Spermatia of corn rust, Puccinia sorghi, 923–925; The spermatia of flax rust, Melampsora lini, 487

Allium, ascalonicum, shallots, resistance to pink root, 294; cepa, resistance to pink root, 292–293; cepa, varietal differences, 295; fistulosum, Nebuka type, 292; fistulosum, resistance to Peronospora schleideni, 109–110; fistulosum, resistance to Phoma terrestris, 109–110; fistulosum, resistance to pink root, 294–297; fistulosum, resistance to Urocystis cepulae, 109–110; porrum, leek, resistance to pink root, 294; sativum, garlic, resistance to pink root, 294; schoenoprasum, chives, resistance to pink root, 294

Alopecurus fulvus, sclerotium from, 867

Alternaria, dianthi, control of serious disease of carnation caused by, 12; rot of citrus in Brazil, 737

Aluminum, -ion, effect on conifer damping-off, 18; sulphate, effect on soil pH, 18

Amaryllis, leaf scorch of, 771

American Phytopathological Society, announcement of meeting, 842; report of Pacific Division, 928; report of the twenty-fourth annual meeting, 489–500

Ananas comosus, host of pineapple mealy bug, 213

Anatomical changes, in big bud of tomato, 648

Anatomy, of "Kroepoek-diseased" leaf of Nicotiana tabacum and of Zinnia elegans, 175–190; of leaves of sugar beet affected by curly top, pathologic changes in, 679–712

ANDERSON, H. W., see THORNBERRY, H. H.

Annona, host of Dothiorella ribis, 929

Anthocyanin, development in intumescences of Eucalyptus coccifera, 285, 288

Anthracnose, of snap-beans in transit, 415–420; use of term as applied to diseases caused by Sphaceloma and Gloeosporium, 389–395

Antirrhinum, inoculated with Verticillium dahliae, 125

Ants, accompanying pineapple mealy bug, 213

Aphanomyces, merely subsidiary in pea wilt, 14; sp., soil fertility in relation to incidence of root rot of peas due to, 36

Aphelenchoides, fragariae, attacking Iris bulbs, 105; unusual symptoms on Chrysanthemum hortorum, 622

Aphids, transmission of some viruses affecting spinach by, 446–474

Aphis, gossypii, vector of bean-mosaic virus, 40; rumicis, vector of bean-mosaic virus, 40

Aplanobacter, stewarti, transmission by larval stage of Diabrotica longicornis, 18

Apple, control of hairy root and crown gall in nursery trees, 30; decay by **Aspergillus sclerotiorum**, n. sp., 306–308; fire blight, weak Bordeaux spray in the control of, 729–733; fruit spot, asso-

ciated with papular type of measles, 4; Gymnosporangium germinale, relation of age of tissue to infection by, 551–552 measles, fruit spot, associated with papular type of, 4; perennial canker, relation of growth cycle and nutrition to, 33; quince rust on, incubation period of, 551; quince rust on, symptoms of, 548–549; resistance to fire blight, 32; rust, caused by Gymnosporangium juniperi-virginianae, 546; scab, development on stored fruit, 5; scab, preventive treatments, effect on tree growth and yield, 11; varietal susceptibility, to quince rust, 547

Appressoria of Stagonospora curtisii, 780

Ardromeda glaucophylla, host of spruce needle rusts in Minnesota, 613, 614

Armillaria mellea, on peach trees in the Carolinas, 28

ARMSTRONG, G. M., and W. B. ALBERT, Downy mildew of tobacco on pepper, tomato, and eggplant, 837–839

Arsenical sprays, zinc hydroxide a substitute for calcium hydroxide in, 17

ARTHUR, J. C., Authority for the name Cronartium ribicola, 559–561; Correction in authority for Cronartium ribicola, 841

Ascochyta citri, causing leaf spot of citrus in Brazil, 736

Ascospore discharge in Ophiobolus graminis, and its probable relation to the development of whiteheads in wheat, 721–728

Aspergillus, niger, cause of citrus decay in Brazil, 737; niger, effect of vitamins on growth of, 929; **sclerotiorum**, n. sp., relation to decay of apples, 306–308

Autoirrigator, improvements in, 928

Bacillus, amylovorus, dissemination by honeybees, 27; amylovorus, resistance of rosaceous plants to, 32; amylovorus, weak Bordeaux spray in the control of, 729–733; aroideae, 26; aroideae, serum-agglutination studies with, 27; carotovorus, 26; carotovorus, serum-agglutination studies with, 27; coli communis, serum-agglutination studies with, 27; coli group, variants of, 27; phytophthorus, 26; phytophthorus, serum-agglutination studies with, 27

Bacteria, seed-borne, main cause of pea wilt, 14; soft-rot, 26, 27; soft-rot, variants of Bacillus coli group, 27; toxicity of pentathionic acid to, 155–174

Bacterial, blight, of broad bean in Louisiana, 27; blight, of snap beans in transit, 420–425; blight, on twig and blossom of raspberry, 21; canker, of Prunus, spp., in California, 36–37; leaf blight, of dent corn, 15–16; speck, of tomatoes, 897; wilt, of corn, 18

Bacterium, atrofaciens, on barley, 10; cucurbitae, on cucumber, 309, 310; marginatum, cause of gladiolus scab, 802; medicaginis var. phaseolicola, on snap beans in transit, 420–425; phaseoli, on snap beans in transit, 420–425; **punctulans**, n. sp., cause of speck of tomato, 87; tabacum, relation of toxin to pathogenicity and host range of, 6; tabacum, serum-agglutination studies with, 27; tumefaciens, causing crown gall on Sahuaro, 21; tumefaciens, on peach trees in the Carolinas, 28; yellow, associated with gum disease of citrus, 734

BAIN, HENRY F., see SHEAR, C. L.

BALD, J. G., see SAMUEL, GEOFFREY

Banana, host of pineapple mealy bug, 213

BANFIELD, W. M., see RIKER, A. J.

Barley, basal glume rot of, 10; root decoction, toxicity to Phymatotrichum omnivorum, 533–534, 536; sclerotium from, 867

BARNES, B. F., see RODENHISER, H. A.

BARSS, H. P., see BRESSMAN, E. N.

BEALE, HELEN PURDY, Serologic reaction as a means of determining the concentration of tobacco-mosaic virus, 4

Bean, mosaic, influence of pollen and ovule infection in seed transmission of, 25; mosaic, transmission of virus by insects, 40; relation of different legume mosaics to, 562–564; rust, nature of resistance to, 38; seedlings, losses caused by Rhizoctonia bataticola, in California, 949–963; species and varieties of, susceptible

to seedling stem blight, in California, 953–957; transmissibility of certain legume-mosaic viruses to, 39
Beans, field, seedling stem blight of, caused by Rhizoctonia bataticola at high temperatures, 949–963
BEAUVERIE, MARIE-ANTOINETTE, Les maladies à ultravirus des plantes, book review of, 926–927
Beech, Nectria coccinea on, 10
Basal glume rot of barley, 10
Basswood, Nectria canker of, 36
Beet, sugar, curly-top leaves, clearing of the veins of, 699; sugar, curly-top leaves, differentiation of xylem in the degenerated bundle cap of, 696; sugar, curly-top leaves, exudate from necrotic phloem of, 698; sugar, curly-top leaves, hypertrophy and hyperplasia in the phloem of, 689; sugar, curly-top leaves, localization of necrosis in the phloem and pericycle of, 686; sugar, curly-top leaves, overgrowths on veins of, 703; sugar, curly top, pathologic changes in anatomy of leaves of, 679–712; sugar, healthy leaves, obliteration in phloem of, 684; sugar, healthy leaves, secondary vascular tissue in the bundle cap of, 694; sugar, healthy leaves, structure of the phloem and pericycle region in, 682
Bentonite, use with insoluble copper compounds in fungicides, 561–562
BERG, ANTHONY, A fruit spot associated with the papular type of apple measles, 4
BERGMAN, H. F., and W. E. TRURAN, An apparent case of transmission of cranberry false blossom through a natural graft, 670–672
Beta vulgaris, and tobacco, comparison of viruses on, 451–458; curly top, pathologic changes in anatomy of leaves of, 679–712
Big bud, virus disease of tomato, 641–653
Biography, ARTHUR JACZEWSKI, 111–116; of HORACE MANN WOOLMAN, 931–933
BIRKELAND, J. M., Electrophoretic studies on purified plant viruses, 4–5; Experiments on acquired immunity in tobacco mosaic and spot necrosis, 5
Black, rot, of coffee, 875; shank, of tobacco, 605–612
Blackberry, inoculated with Verticillium dahliae, 125
Blister, rust, of Pinus monticola, 917; rusts, hosts of Tuberculina maxima, 299
BLODGETT, F. M., E. O. MADER, O. D. BURKE, and R. B. MCCORMACK, Total amount of copper applied and ratio of lime to copper in Bordeaux as important factors in potato spraying, 5
BLOOD, H. L., see RICHARDS, L. A.; see WANN, F. B.; B. L. RICHARDS, and F. B. WANN, Studies of psyllid yellows of tomato, 930
Bog rosemary, host of spruce needle rusts in Minnesota, 613, 614
Book review, Fungous diseases of the cultivated cranberry, 314–315; Insects and diseases of ornamental trees and shrubs, 312–314; Les maladies à ultravirus des plantes, 926–927; Phytopathological and botanical research methods, 1004–1005; Principles of plant physiology, 1005–1006; Root-nodule bacteria and leguminous plants, 625–626
Bordeaux, in potato spraying, 5; mixture, evidence of stimulation of potatoes by, 22; mixture, for control of gladiolus scab, 812; mixture, in control of coffee black rot, 885; mixture, notes on, 31–32; mixture, relation of, to potato tip burn, 916; spray, weak, in the control of fire blight of apple, 729–733
Botrytis, cinerea, cause of potato tuber rot, 11, 993–999; cinerea, on snap beans in transit, 436–437; sp., cause of stem canker of China aster, 20; spp., on narcissus, 773
BOYD, O. C., see GILGUT, C. J.
BRATLEY, C. O., Development of apple scab on stored fruit, 5
Bremia lactucae, physiologic forms on lettuce, 18–19
BRENTZEL, W. E., Physiologic specialization of Tilletia tritici on emmer, 483–485
BRESSMAN, E. N., and H. P. BARSS, Experiments with head smut of corn in western Oregon, 396–403; and R. A. NICHOLS, Germination of the oospores of Pseudoperonospora humuli, 485–487
Brevicoryne brassicae, vector of bean-mosaic virus, 40

BRIERLEY, PHILIP, Virus diseases of dahlia, 6
Broad bean, bacterial blight of, 27
BROADFOOT, W. C., On the pathogenicity of Wojnowicia graminis, 1001–1002
BROOKS, A. N., A wilt of strawberry caused by Colletotrichum fragariae, 6
Brooming disease, of black locust, transmissibility of, 83–90; of Robinia pseudoacacia transmitted by grafts, 13
BROWN, A. M., see NEWTON, MARGARET
BROWN, J. G., and M. M. EVANS, Crown gall on a conifer, 97–101
Brown, patch, caused by Sclerotium rhizodes, 8; patch, of turf, effect of temperature on, 8; rot, of peach, distribution of losses from, in the United States, 978; rot, of peach, in cars, in the United States, 980
BRYAN, MARY K., Bacterial speck of tomatoes, 897–904; Bacterium cucurbitae on cucumber, 309
BUHRER, EDNA M., see STEINER, G.
Bunt, of wheat, average yearly losses from, in the United States, 977; of wheat, in cars, percentage of, 978; of wheat, losses from, in the United States, 976
Bureau of Crop Estimates, estimating losses from plant diseases, method of, 976
BURKE, O. D., see BLODGETT, F. M.
BUSCH, HANS J., and FREDERICK A. WOLF, Manufactured tobacco, a source of inoculum for mosaic in flue-cured tobacco, 839–841
BUTLER, WALLACE, see STAKMAN, E. C.

Calochlor, treatment with, for control of gladiolus scab, 807, 810, 811
Calogreen, treatment with, for control of gladiolus scab, 805, 806, 810, 811
Camarosporium, associated with brooming disease of Robinia pseudoacacia, 13
Canadian Phytopathological Society, advance notice of meeting, 410
Canker, of stem of Crotalaria spectabilis caused by **Diaporthe crotalariae**, n. sp., 596–604; on hollyhock stem caused by Sclerotinia libertiana, 39; records of, on Citrus from herbarium specimens, 820
Cankers, of China aster stems, 20; of peach, isolations of Phytomonas pruni from, 794; spring, of P. pruni, description of, on peach, 787; summer, of P. pruni, description of, on peach, 787
Carbohydrates, utilization by Phytophthora nicotianae in cultures, 611
Carbon, dioxide, effect on cultivated mushrooms, 20; disulphide emulsion, control of root-knot nematode in greenhouses with, 13; source of, effect on sporulation of Phyllosticta solitaria, 518–519
Carnation, control of serious disease caused by Alternaria dianthi, 12
Carnegiea gigantea, crown gall on, 21
Carrisia, host of Dothiorella ribis, 929
CARTER, WALTER, The pineapple mealy bug, Pseudococcus brevipes, and wilt of pineapples, 207–242; The spotting of pineapple leaves caused by Pseudococcus brevipes, the pineapple mealy bug, 243–259
Carya pecan, host of Dothiorella ribis, 929
Cedar, red, susceptibility to apple rust, 546–547; red, susceptibility to hawthorn rust, 546–547; red, susceptibility to quince rust, 546–547
Celery, breeding for resistance to Fusarium yellows, 25
Centrosome, existence in Puccinia malvacearum, 581
Cephalosporium, lecanii, in Brazil, 737; on elm, 21
Ceratonia, host of Dothiorella ribis, 929
Cercospora, arachidicola Hori, leaf spot of peanut, 634; arachidis P. Henn., synonym of C. personata (B. & C.) E. & E., 637; arachidis P. Henn. var. macrospora Maff., synonym of C. arachidicola Hori, 634; personata (B. & C.) E. & E., leaf spot of peanut, 637
Cercosporella herpotrichoides, occurrence in a specific ecological area on wheat, 33–34
Cereal diseases, method of estimating loss of yield from, 12
Cereals, foot rot, relation of mechanical injury to, 22
Ceresan for control of oat smut, 827, 829
Chamaedaphne calyculata, host of Melampsoropsis cassandrae, 615

CHANDLER, NORMAN, see JAGGER, IVAN C.
CHAPMAN, A. DALE, see LINDGREN, RALPH M.
CHARLES, VERA K., see MEIER, F. C.
China aster, Fusarium strains in relation to wilt in, 29–30; two stem cankers of, 20
Chionodoxa luciliae, infested with Tylenchus dipsaci, 620
Chives, Allium schoenoprasum, resistance to pink root, 294
Chlamydospores of Tolyposporium filiferum, germination of, 987–989
Chloris gayana, host of pineapple mealy bug, 249
Chlorosis in suspected mosaic of strawberry, 656
Chromosome, number in Puccinia malvacearum, 581
Chrysanthemum hortorum, Aphelenchoides fragariae on, 622
Citrus, algal leaf spot, 736; Alternaria rot, 737; anthracnose, 735–736; Aspergillus rot, 737; black melanose, 736; blue and green mold, 737; damping off, 735; diseases, observations and notes in Minas Geraes, Brazil, 734–737; entomogenous fungi, 737; felt 736; Fusarium rot, 737; gum disease, 734; gum disease, Fusarium isolated from, 734; gum disease, Phytophthora isolated from, 734; gum disease, yellow bacterium associated with, 734; leaf spotting, 736; melanose, 735; new locations for P. citrophthora and P. hibernalis, 667–669; Penicillium digitatum on, 737; Penicillium italicum on, 737; Phytophthora citrophthora on, in Florida and Louisiana, 667–669; P. hibernalis on, in California, 667–668; P. parasitica on, in Florida, 667, 668; psorosis, a virus disease of, 930; records of canker of, from herbarium specimens, 820; root rot, 734–735; root rot, Fusarium associated with, 734; root rot, Nectria associated with, 735; scab, 735; scab, common, distinct from sweet-orange scab in Argentina, 479; scab, distribution of, 481; scab, records of, from herbarium specimens of the genus Citrus in England and the United States, 475–482; scaly bark of, 930; sinensis, navel orange, Sphaceloma fawcettii var. **viscosa**, n. var., on, in Brazil, 538–545; sooty mold, 736; sour rot, 737; spp., hosts of Dothiorella ribis, 929
Cladosporium personatum B. & C., synonym of Cercospora personata (B. & C.) E. & E., 637
CLAYTON, E. E., Relation of the toxin produced by Bacterium tabacum to the pathogenicity and host range of this organism, 6
Clover, potato-leaf-hopper injury, 19; red, ringspot-like virus disease of, 746–747
COCHRAN, L. C., see NELSON, RAY
Coffee, black rot of, in Mysore, 875
Colchicum speciosum album, host of Tylenchus dipsaci, 620
Colletotrichum, falcatum, physiologic forms of, 557–559; fragariae, a wilt of strawberry caused by, 6; gloeosporioides, on citrus in Brazil, 735–736; lindemuthianum, on snap beans in transit, 415–420
Colloidal copper, a homemade spray of, 29
Comparison of pycnial stage of Cronartium ribicola on Pinus lambertiana and P. monticola, 204–205
Concentration of virus, 752
CONDIT, IRA J., and W. T. HORNE, Mosaic of fig in California, 887–896; A mosaic of the fig in California, 7
Conifers, damping off, 18
Control, of fire blight, apple, by weak Bordeaux spray, 729–733; of gladiolus scab, 802; of oat smut, 825; of olive knot, 37; of powdery mildew of crape myrtle, 818; of tobacco mosaic, 831
COOK, HAROLD T., Infection of seed clusters of spinach by Peronospora effusa, 7
COOK, MELVILLE T., Pineapple leaf spot, 7
COOLEY, J. S., see SHEAR, ELMER V.
Copper, amount applied and ratio to lime in Bordeaux for potato spraying, 5; compounds, insoluble, for fungicides, 561–562; -lime dust, for control of gladiolus scab, 812; sulphate, form of toxicity surface in relation to conidia of Sclerotinia americana, 23
Cork-borer method for inoculating trees, 487–488
Corm treatment for gladiolus scab, 803

Corn, bacterial leaf blight of, 15–16; bacterial wilt, insect transmission of, 18; bacterial wilt of, 18; bacterial wilt, relation of various bacterial cultures to, 18; bound water in leaf tissues of heat-resistant and heat-susceptible inbred strains of yellow-dent, 15; Gibberella moniliformis on, 368; head smut, in western Oregon, 396–403; Indian, a host of pokkah-bong, 672; Physalospora zeicola on, 63–72; reaction of seedlings of, to Gibberella saubinetii, 905; rust, spermatia of, 923; seedling blight, nature of resistance to, 9

Corona, oat dust, for control of smut, 827, 829; 174 E, for control of oat smut, 827

Correction to article (Macrophomina phaseoli and Rhizoctonia bataticola); by W. W. MACKIE, 205

Corticium koleroga, cause of coffee black rot, 875; hyphal fusions in, 878, 879

Cotton, inoculated with Verticillium dahliae, 125; **Ozonium texanum**, n. sp., on stalks and roots of, 24; root decoction, growth of Phymatotrichum omnivorum on, 529–533, 535; root decoction, toxicity to P. omnivorum, 529–533, 535; -root-rot fungus, fusion of hyphae, 676; -root-rot fungus, morphology and life history of, 24

Cottony leak on snap beans in transit, 428

Cowpea, blackeye, resistance to Rhizoctonia bataticola, 953

Cranberry, production in New Jersey, 984; production in New Jersey, effect of false blossom on, 984; relation of varieties to spread of false blossom, 36; transmission of false blossom through a natural graft, 670

Crape myrtle, powdery mildew of, 1002–1003

Crataegus, Arnoldiana, Gymnosporangium germinale on, in New York, 546; beata, G. germinale on, in New York, 546; brainerdi, G. germinale on, in New York, 546; calpodendron, G. germinale on, in New York, 546; filipes, G. germinale on, in New York, 546; G. germinale, relation of age of tissue to infection by, 551–552; holmesiana, G. germinale on, in New York, 546; holmesiana, G. germinale on, perennial infection by, 549; monogyna, G. germinale on, in New York, 546; oxyacantha, G. germinale on, in New York, 546; quince rust on, incubation period of, 551; quince rust on, symptoms of, 548

CREAGER, DON B., Fusarium basal rot of bulbous iris, 7; Leaf scorch of narcissus, 770–786

Crinkle disease, strawberry, transmission of, 738–740

Cronartium, cerebrum, host of Tuberculina maxima, 299; coleosporioides, host of T. maxima, 299; commandrae, host of T. maxima, 299; comptoniae, host of T. maxima, 299; harknessii, host of T. maxima, 299; ribicola, authority for name, 203–204, 559–561; ribicola, comparison of pycnial stage on Pinus lambertiana and P. monticola, 204–205; ribicola, correction in authority for, 841; ribicola, field inoculations of Pinus strobus with sporidia of, 105–107; ribicola, Fischer and C. ribicola Dietrich, 203–204; ribicola, host of T. maxima, 299; ribicola, resistance to, of current season's shoots of P. monticola, 917; stalactiforme, host of T. maxima, 299

CROSIER, WILLARD, Culture of Phytophthora infestans, 713–720

Crotalaria spectabilis, stem canker of, by **Diaporthe crotalariae**, n. sp., 596–604

Crown gall, control in unions of piece-root-grafted apple trees, 30; on Cupressus arizonica, 97–101; on Sahuaro (Carnegiea gigantea), 21; seasonal development on apple trees in nursery, 29

Cryptococcus fagi, relation to beech disease caused by Nectria coccinea, 10

Cuban streak of sugarcane, 674

Cucumber, and spinach, comparison of viruses on, 466–467; Bacterium cucurbitae on, in Massachusetts, 309, 310; inoculated with Verticillium dahliae, 117–154; mosaic, inheritability of virus in aphids, 467–468; mosaic, natural infection of spinach, 466; mosaic, on tobacco, 311; mosaic, transmission to spinach, 459–463

Cucumis sativa and spinach, comparison of viruses on, 466–467
Culture, media, for growth studies of Phymatotrichum omnivorum, 527, 529; media, influence on sporulation of Phyllosticta solitaria, 505–519; medium, humus-extract agar for oospore production in Pythium, 745; medium, K. U., for Phyllosticta solitaria, 514; of Phytophthora infestans, 713–720
Cultures, of Diaporthe crotalariae, 600; of Tolyposporium filiferum, 990–992; on no-nitrogen medium, of Helminthosporium sativum, 617
Cupressus arizonica, crown gall on, 97–101
Curly top, differentiated from psyllid yellows, 930; of tomato, chemical changes accompanying, 929; sugar beet, pathologic changes in anatomy of leaves, 679–712
Cydonia oblonga, host of Dothiorella ribis, 929
Cytological investigations on spike disease of sandal, Santalum album, 191–202
Cytology of teliospores, promycelia, and sporidia in Puccinia malvacearum, 572–586
Cytospora canker of Picea spp., 11
Czapek's agar as a medium for growth of Helminthosporium sativum, 617

Dactylis glomerata, sclerotium from, 867
DAHL, ARNOLD S., Effect of temperature on brown patch of turf, 8
Dahlia, inoculated with Verticillium dahliae, 117–154; mosaic, 6; ring spot, 6; virus diseases of, 6
Damping-off, effect of H-ion and aluminum-ion on destructiveness in conifers, 18; of spinach combated by seed treatment, 28; use of formaldehyde dust to control, 38
Datura stramonium, affected by mites, 622
DAVIS, W. H., Snow mold and brown patch caused by Sclerotium rhizodes, 8
Diabrotica longicornis, transmission of corn bacterial wilt by larval stage of, 18
Diaporthe crotalariae, n. sp., isolation of, 599, 600; occurrence of, 596, 597; pathogenicity of, 601; taxonomy of, 602; technical description, 602
DICKINSON, SYDNEY, The technique of isolation in microbiology, 357
DICKSON, A. D., see DICKSON, JAMES G.
DICKSON, JAMES G., and R. G. SHANDS, Disease-resistant wheats recently introduced from Russia, 8–9; KARL P. LINK, and A. D. DICKSON, Nature of resistance of corn to seedling blight, 9
Die-back of twigs, caused by Phytomonas pruni, 790
DIETRICH, preferable authority for Cronartium ribicola, 203–204
Digitalis purpurea, host of Tylenchus dipsaci, 620
Diospyros, host of Dothiorella ribis, 929
Diplodia, frumenti, relation to Physalospora zeicola on corn, 63–68; tubericola, hosts of, 70–72
DIPPENAAR, B. J., Environment and seed treatment in relation to common potato scab, 9
Disease-resistant wheats introduced from Russia, 8–9
Diseases, environmental factors in relation to snap bean, occurring in shipment, 411–445; sclerotium, of grains and grasses, 863
DOOLITTLE, S. P., review of Les maladies à ultravirus des plantes, by MARIE-ANTOINETTE BEAUVERIE, 926–927
DOPP, ERNEST, see RANDS, R. D.
Dothiorella ribis, inoculations with, 929
DOWN, E. E., see NELSON, RAY
Downy mildew, germination of oospores of, on hops, 485–487; of tobacco on pepper, tomato, and eggplant, 837
DRAYTON, F. L., Perfect stage of Sclerotium gladioli, 9
Dubay, 500 LL, for control of oat smut, 827, 829; 952 C, for control of oat smut, 827, 829; 952 K, for control of oat smut, 827, 829; 1134, for control of oat smut, 829
DUFRÉNOY, J., Differentiation of green- and yellow-mosiac viruses in tobacco, 10
DUGGAR, B. M., and BURT JOHNSON, Stomatal infection with the virus of typical tobacco mosaic, 934–948
Dusting for control of gladiolus scab, 812

EARDLEY, C. M., see SAMUEL, GEOFFREY
Ecology of Cercosporella foot rot of wheat, 33–34
EDDINS, A. H., and R. K. VOORHEES, Physalospora zeicola on corn and its taxonomic and host relationships, 63–72
Edge wilt, synonym of pineapple wilt, 208
Eggplant, downy mildew of tobacco on, 837; susceptibility of, to Verticillium albo-atrum, 125; susceptibility of, to V. dahliae, 125; V. hadromycosis, 130–138
EHRLICH, JOHN, Nectria coccinea on beech, 10
Electrophoresis on purified plant viruses, 4–5
ELLIOTT, CHARLOTTE, see HOLBERT, J. R.; and A. G. JOHNSON, Basal glume rot of barley, 10
Elm, Cephalosporium on, 21; diseases in America, 21; Dutch elm disease, in America, 21; root disease of, 21; Verticillium on, 21
Elsinoë canavaliae Rac., cause of Lima-bean scab, 662
Emmer, physiologic specialization of Tilletia tritici on, 483–485
Environal studies on Verticillium hadromycosis, 117–154
Environment in relation to common potato scab, 9
Environmental, factors in relation to snap-bean diseases occurring in shipment, 411–445; relations of Heterodera radicicola, 41–62
Epiphytotics of seedling stem blight of bean in California, 949–953
Eragrostis aspera Nees, host of **Sclerospora butleri**, n. sp., 587–595
Eriobotrya japonica, host of Dothiorella ribis, 929
Eriophyes fici, mites of fig, 893
Erysiphe, graminis, resistant wheats from Russia, 8; lagerstroemiae, comparison with Phyllactinia corylea, on crape myrtle, 1002–1003; **lagerstroemiae**, n. sp., on crape myrtle, 814; scandens, synonym of Sclerotium koleroga, 876
ESAU, KATHERINE, Pathologic changes in the anatomy of leaves of the sugar beet, Beta vulgaris L., affected by curly top, 679–712
Etiological, classification of control measures, 861; classification of pathic events, 860; phytopathology, 843
Eucalyptus, coccifera, intumescences on, 284; cornuta, intumescences on, 281; host of Dothiorella ribis, 929

Fagus grandifolia, Nectria coccinea on, 10
False blossom, of cranberry, 670–672; of cranberry, in New Jersey, effect on production, 984; of cranberry, transmission through natural graft, 670; relation of cranberry varieties to spread of, 36
FARIS, J. A., Influence of soil moisture and soil temperature on infection of wheat by Urocystis tritici, 10–11
FAWCETT, HOWARD S., New locations for Phytophthora citrophthora and P. hibernalis on Citrus, 667–669; New symptoms of psorosis, indicating a virus disease of Citrus, 930; and ANNA E. JENKIN . Records of citrus canker from herbarium specimens of the genus Citrus in England and the United States, 820–824
Feijoa, host of Dothiorella ribis, 929
FELIX, E. L., Disease resistance in Allium fistulosum, 109–110
FELT, E. P., and W. H. RANKIN, Insects and diseases of ornamental trees and shrubs, review of, 312–314
Ficus, mosaic of, 891
Fig, mites, 893; mosaic in California, 7; mosaic of, 887; scale, Mediterranean, in California, 894; Stilbum cinnabarinum the cause of a new disease of, 35
Filtration, effect of, on thermal death point of virus, 754
Fire of narcissus, 773
Fire blight, apple, weak Bordeaux spray in the control of, 729–733; resistance of rosaceous plants to, 32–33
Flax, spermatia of rust of, 487
FOLSOM, DONALD, Botrytis cinerea as a cause of potato tuber rot, 993–999; Effect of scab-preventive treatments on apple-tree growth and yield, 11; Potato tuber rot caused by Botrytis cinerea, 11
Food, quality, influence on sporulation of Phyllosticta solitaria, 511–515; quantity, influence on sporulation of P. solitaria, 506–515

Foot rot, relation in cereals to mechanical injury, 22
Forest products, damage by fungi to, 13; in the United States, losses from decay of, 983
Formaldehyde, dust, use with vegetable seedlings, 38; effect on vitality of potato sets, 277; for control of gladiolus scab, 804; for control of oat smut, 829; toxicity to sclerotia of Rhizoctonia solani, 272–273
Foxtail, sclerotium from, 867
Fraxinus, host of Dothiorella ribis, 929; nigra, Nectria canker of, 36
FRED, E. B., I. L. BALDWIN, and ELIZABETH MCCOY, Root nodule bacteria and leguminous plants, book review of, 625–626
FRYE, B. C., see HOLBERT, J. R.
Fungi, as affected by vitamins, 929; damage to forest products by, 13; toxicity of pentathionic acid to, 155–174
Fungicides, promising, 561–562
Fusarium, associated with damping-off of citrus, 735; associated with gum disease of citrus, 734; associated with root rot of citrus, 734; basal rot of bulbous iris, 7; conglutinans var. callistephi, pathogenic to China aster, 29–30; conglutinans var. majus, pathogenic to China aster, 30; culmorum var. letius, cause of foot rot of cereals, 22; lateritium var. fructigenum, pathogenic to China aster, 30; merely subsidiary in pea wilt, 14; niveum, relation to formation of tyloses in melon plants, 33; orthoceras var. pisi, comparative studies of peas resistant and susceptible to wilt caused by, 36; oxysporum f. 6, pathogenic to China aster, 30; rot of citrus, in Brazil, 737; sp., cause of celery yellows, 25; strains in relation to wilt in China aster, 29–30; wilt, of muskmelon, in Minnesota, 554–556
Fusion of hyphae of the cotton-root-rot fungus, 676

Garlic, Allium sativum, resistance to pink root, 294
GARRETT, S. D., see SAMUEL, GEOFFREY
Gelatin, effect of, on tobacco-mosaic virus, 756
Germination, of oospores of Pseudoperonospora humuli, 485–487; of spores of Gymnosporangium germinale, 539–551
Gibberella, moniliformis, on corn, 368; saubinetii, reaction of maize seedlings to, 905
GILGUT, C. J., and O. C. BOYD, Cytospora canker of Picea spp., 11
Girdling of Persian (English) walnuts grafted on black-walnut stocks, 408–409
Gladiolus, scab, control of, 802; scab, control of, by corm treatment, 803; scab, control of, by spraying and dusting, 812; **Sclerotinia gladioli**, n. comb., the perfect stage of Sclerotium gladioli on, 9
Gloeosporium, diseases caused by, "anthracnose" and "scab" applied to, 389–395; perennans, relation of growth cycle and nutrition of apple to infection by, 33
GODFREY, G. H., see HOSHINO, HELENE MORITA; and HELENE MORITA HOSHINO, Studies on certain environmental relations of the root-knot nematode, Heterodera radicicola, 41–62
Gold Dust, treatment with, for control of gladiolus scab, 808
GOLDSWORTHY, M. C., and E. L. GREENE, Some promising fungicides, 561–562
Graft, disorder, of Persian (English) walnuts on black-walnut stock, 408–409; natural, as a means of transmission of false blossom of cranberry, 670; transmission of big bud in tomato, 650
Grains, Sclerotium diseases of, 17, 863
Grapefruit, Nematospora coryli on, in Florida, 384
Grasses, Sclerotium diseases of, 17, 863
Gray-mold rot on snap beans in transit, 436–437
GREANEY, F. J., Method of estimating loss in yield from cereal diseases, 12; see MACHECEK, J. E.; and J. E. MACHACEK, Production of a white fertile saltant of Helminthosporium sativum by means of ultra-violet radiation, 12; and J. E. MACHACEK, The production of a white fertile saltant of Helminthosporium

sativum by means of ultra-violet radiation, 379
Green spotting of pineapple leaves, 243; relation of mealy bug to, 243–246; anatomy of, 248; transmission experiments, 253
GREENE, E. L., see GOLDSWORTHY, M. C.
Growth, in culture, of Phymatotrichum omnivorum, 525–537
GUBA, E. F., Control of Alternaria dianthi causing a serious disease of the carnation, 12; Control of the root-knot nematode in greenhouses with carbon disulphide emulsion, 13; "Suspected mosaic" of strawberry, 654–661
GUNN, K. C., see NEAL, D. C.
Gymnosporangium, germinale, 546–553; germinale, causing witches' brooms on red cedar, 547; germinale, in galls of G. globosum, 547; germinale, incubation period of, 551; germinale, on Crataegus holmesiana, perennial infection, 549; germinale, on Crataegus spp., in New York, 546; germinale, pathogenic strains of, 549; germinale, relation of age of host tissue to infection by, 551–552; germinale, spore germination, 549–551; germinale, susceptibility of apple varieties to, 547; germinale, susceptibility of red cedar to, 546–547; germinale, symptoms, on apple, 548–549; germinale, symptoms, on Crataegus, 548; globosum, galls preempted by G. germinale, 547; globosum, susceptibility of red cedar to, 546–547; juniperi-virginianae, susceptibility of red cedar to, 546–547; nidus-avis, 547

Hairy root, control in unions of piece-root-grafted apple trees, 30; seasonal development on apple trees in nursery, 29
Halo of Cercospora arachidicola on peanut, 634
HARTER, L. L., book review of Phytopathological and botanical research methods, by THOMAS ELLSWORTH RAWLINS, 1004–1005; see LAURITZEN, J. I.
HARTLEY, CARL, Damage to forest products by fungi, 13; see JACKSON, L. W. R.; and L. W. R. JACKSON, A brooming disease of Robinia pseudoacacia transmitted by grafts, 13
HATCH, A. B., Pure-culture technique for quantitative studies of plant growth in association with microorganisms, 14; True mycorhizal fungus in contrast to Mycelium radicis atrovirens, 14
Hawthorn rust, caused by Gymnosporangium globosum, 546
HAYES, H. K., I. J. JOHNSON, and E. C. STAKMAN, Reaction of maize seedlings to Gibberella saubinetii, 905–911
HEIMBECK, LOUISE SOLBERG, Seed-borne bacteria main cause of pea wilt; Fusarium, Aphanomyces, and other organisms merely subsidiary, 14
Helminthosporium, sativum, effect of vitamins on growth of, 929; sativum, mutation in a no-nitrogen medium, 617–619; sativum, production of white fertile saltant by means of ultra-violet radiation, 12, 379; sigmoideum, genetic connection with Sclerotium oryzae and Leptosphaeria salvinii, 35
HENDERSON, R. G., Increasing the resistance of tobacco ring-spot virus to aging in vitro by use of carbolic acid, 14–15
Hendersonia curtisii, synonym for Stagonospora curtisii, 773
Herbarium specimens, records of citrus scab from, in the genus Citrus in England and in the United States, 475–482
Heterodera radicicola, control of in greenhouses, 13; on peach trees in the Carolinas, 28; studies on certain environmental relations of, 41–62; thermal death point, 260–270
Heterothallism, in Melampsora lini, 4; in Puccinia coronata, 4; in P. graminis, 4
HEUBERGER, J. W., see NORTON, J. B. S.; and J. B. S. NORTON, Water loss in tomato mosaic, 15
Hicoria spp., anther blight, 407
Hieracium venosum, intumescences on, 285
HILDEBRAND, E. M., see RIKER, A. J.
HINES, LEE, see STAKMAN, E. C.
Hippeastrum vittatum, leaf scorch of hybrids, 771
HOGGAN, ISMÉ A., Some viruses affecting spinach, and certain aspects of insect transmission, 446–474

HOLBERT, J. R., and B. C. FRYE, Bound water in leaf tissues of heat-resistant and heat-susceptible inbred strains of yellow-dent corn, 15; CHARLOTTE ELLIOTT, and BENJAMIN KOEHLER, Bacterial leaf blight of dent corn, 15–16

Hollyhock, cytology of Puccinia malvacearum on, 572–586; Sclerotinia stem canker of, 39

HOLTON, C. S., Differences in size and shape of smut balls of two physiologic forms of Tilletia tritici, 16; Inheritance of chlamydospore characteristics in oat-smut fungi, 16

Honeybees, dissemination of Bacillus amylovorus by, 27

Hops, germination of oospores of downy mildew of, 485–487

HORNE, W. T., see CONDIT, IRA J.

HORSFALL, JAMES G., and Z. I. KERTESZ, Influence of root rot on peas; March of the disease, 16–17

HOSHINO, HELENE MORITA, see GODFREY, G. H.; and G. H. GODFREY, Thermal death point of Heterodera radicicola in relation to time, 260–270

HUBER, G. A., **Aspergillus sclerotiorum,** n. sp., and its relation to decay of apples, 306–308

HUMPHREY, HARRY B., biography of HORACE MANN WOOLMAN, 931–933

Humus-extract agar for oospore production in Pythium, 745

HUNGERFORD, C. W., see REMSBERG, RUTH; and RUTH REMSBERG, Sclerotium diseases of grains and grasses, 17

HURT, R. H., Zinc hydroxide, a substitute for calcium hydroxide in arsenical sprays, 17

HUTCHINS, LEE M., Peach mosaic, 17

Hyalopterus atriplicis, vector of bean-mosaic virus, 40

Hydrogen-ion, concentration, effect on sporulation of Phyllosticta solitaria, 519; effect on conifer damping-off, 18

Hypertrophy and hyperplasia in the phloem of curly-top-affected sugar-beet leaves, 689

Hyphal fusions in Corticium koleroga, 878, 879

Illinoia, pisi, vector of bean-mosaic virus, 40; solanifolii, vector of bean-mosaic virus, 40

Immunity, acquired, in tobacco mosaic, 5; acquired, in tobacco spot necrosis, 5; of figs, from mosaic, 892

Incubation period of Gymnosporangium germinale, 551

Infection, discussion of, 854

Inheritability of cucumber-mosaic virus by progeny of infective aphids, 467–468

Inoculation, cork-borer method for trees, 487–488

Insect transmission, certain aspects of, with some viruses affecting spinach, 446–474; of bacterial wilt of corn, 18

Insects, toxicity of pentathionic acid and salts to, 155–174; transmission of bean-mosaic virus by, 40

Intracellular bodies in tobacco ring spot, 38

Intumescences on, Eucalyptus coccifera, 284; E. cornuta, 281; Hieracium venosum, 285; Mitchella repens, 286; Thurberia thespesioides, 287

Iodine for control of oat smut, 829

Iris, bulbous, Aphelenchoides fragariae on, 105; Fusarium basal rot of, 7

Irises attacked by Tylenchus dipsaci, 103–105

Isolation, single cells, in microbiology, 357

IVANOFF, S. S., Bacterial wilt of corn, 18

JACKSON, L. W. R., Effect of H-ion and aluminum-ion concentrations on conifer damping-off, 18; Effect of sulphuric acid and aluminum sulphate, as used for the control of damping-off of conifers, on soil pH, 18; see HARTLEY, CARL; and CARL HARTLEY, Transmissibility of the brooming disease of black locust, 83–90

JACZEWSKI, ARTHUR, biography of, 111–116; memorial, 501

JAGGER, IVAN C., and NORMAN CHANDLER, Physiologic forms of Bremia lactucae on lettuce, 18–19

JENKINS, ANNA E., Application of the terms "anthracnose" and "scab" to plant diseases caused by Sphaceloma and Gloeosporium, 389–395; Further studies of Lima-bean scab, 662–666; see FAW-

CETT, HOWARD S.; see MASSEY, L. M.; A Sphaceloma attacking Navel orange from Brazil, 538–545; and H. S. FAWCETT, Records of citrus scab mainly from herbarium specimens of the genus Citrus in England and the United States, 475–482

JENSEN, JAMES H., Isolation of yellow-mosaic viruses from plants infected with tobacco mosaic, 964–974

JOHNSON, A. G., see ELLIOTT, CHARLOTTE

JOHNSON, BURT, see DUGGAR, B. M.

JOHNSON, E. M., A ringspot-like virus disease of red clover, 746–747

JOHNSON, HOWARD W., Nature of injury caused by potato leaf hopper on forage legumes, 19

JOHNSON, I. J., see HAYES, H. K.

JOHNSON, JAMES, Cucumber mosaic on tobacco in Wisconsin (title only), 19; Cucumber mosaic on tobacco in Wisconsin, 311

JOHNSON, T., see NEWTON, MARGARET

JOHNSTON, C. O., and E. C. MILLER, Effect of leaf-rust infection on water economy and growth of two wheat varieties, 19–20

JONES, H. A., see PORTER, D. R.

JONES, L. R., biography of Arthur Jaczewski, 111–116; see RIKER, REGINA S.; and REGINA S. RIKER, Two stem cankers of the China aster, 20

Juglandaceae, bacterial (?) blight of, 407–408; new anther blight of, 407–408

Juglans, graft disorder of, girdling, 408–409; regia, Dothiorella ribis from, 929; spp., anther blight, 407; spp., hosts of Dothiorella ribis, 929

Juniperus virginiana, susceptibility to Gymnosporangium spp., 546–547; witches' brooms, caused by G. germinale, 547

KAMAT, M. N., Observations on Tolyposporium filiferum, cause of "long smut" of sorghum, 985–992

KEITT, G. W., see RIKER, A. J.

KENDRICK, JAMES B., Seedling stem blight of field beans caused by Rhizoctonia bataticola at high temperatures, 949–963

KERLING, L. C. P., Anatomy of "Kroepoek-diseased" leaf of Nicotiana tabacum and of Zinnia elegans, 175–190

KERTESZ, Z. I., see HORSFALL, JAMES G.

KEUR, JOHN Y., Seed transmission of the virus causing variegation of Abutilon, 20

KOCH, KARL LEE, The nature of potato rugose mosaic, 319–342

KOEHLER, BENJAMIN, see HOLBERT, J. R.

Koleroga, or black rot of coffee, 873

Kroepoek, a virus disease of tobacco, 175; virus disease of zinnia, 175

Labrador tea, host of Melampsoropsis abietina, 615; host of M. ledicola, 615

LACHMUND, H. G., Resistance of the current season's shoots of Pinus monticola to infection by Cronartium ribicola, 917–922

Lagerstroemia, indica, powdery mildew of, 814, 818; parviflora, powdery mildew of, 814

LAMB, HOWARD, see PIERSTORFF, A. L.

LAMBERT, E. B., Effect of carbon dioxide on cultivated mushrooms, 20

Land-utilization program, contribution of plant pathology to, 404–406

LA RUE, CARL D., Intumescences on leaves of Eucalyptus cornuta, E. coccifera, Hieracium venosum, Mitchella repens, and Thurberia thespesioides, 281–289

Lathyrus, odoratus, mosaic, 562

LAURITZEN, J. I., L. L. HARTER, and W. A. WHITNEY, Environmental factors in relation to snap-bean diseases occurring in shipment, 411–445

LEACH, J. G., A destructive Fusarium wilt of muskmelons, 554–556

Leaf, rust, effect of infection on water economy and growth of two wheat varieties, 19–20; scorch, of narcissus, 770; scorch, of narcissus, symptoms of, 771; spot, of peanut, 627; spot, of pineapple, 7

Leather leaf, host of Melampsoropsis cassandrae, 615

Leaves, pathologic changes in anatomy of sugar beet affected by curly top, 679–712

LEDINGHAM, G. A., Life history, morphology, and cytology of Polymyxa graminis, 20–21

Ledum groenlandicum, host of Melampso-

ropsis abietina, 615; host of M. ledicola, 615
Leek, Allium porrum, resistance to pink root, 294
Legume, mosaic, transmissibility of viruses to bean, 39; mosaics, in relation to beans and peas, 562–564
Legumes, nature of injury caused by potato leaf hopper on, 19
LEHMAN, S. G., Bacterial twig and blossom blight of raspberry, 21
Lentinus tigrinis, growth as affected by temperatures, 73–81
Lenzites sepiaria, growth as affected by temperature, 73–81
LEONARD, LEWIS T., review of root nodule bacteria and leguminous plants, by E. B. FRED, I. L. BALDWIN, and ELIZABETH MCCOY, 625–626
Leprosis of citrus, possible virus origin of, 930
Leptosphaeria salvinii, ascigerous stage of Sclerotium oryzae, 35
Lettuce, physiologic forms of Bremia lactucae on, 18–19
LEVINE, MICHAEL, Crown gall on Sahuaro (Carnegiea gigantea), 21
Life history of Stagonospora curtisii, 775
Light, influence on sporulation of Phyllosticta solitaria, 504–505
Lilac fungus, behavior in British Columbia, 301–302; range in British Columbia, 300; relation to control of white-pine blister rust, 302–304; Tuberculina maxima, hosts of, 299
Lima-bean, scab, 662–666
Lime, treatment with, for control of gladiolus scab, 810, 811
LIMING, O. NEAL, Elm diseases in America, 21; Preparation and properties of pentathionic acid and its salts; its toxicity to fungi, bacteria, and insects, 155–174
LINDGREN, RALPH M., Decay of wood and growth of some Hymenomycetes as affected by temperature, 73–81; Field observations of needle rusts of spruce in Minnesota, 613–616; and A. DALE CHAPMAN, Field inoculations of Pinus strobus with sporidia of Cronartium ribicola in Minnesota, 105–107
LINK, GEORGE K. K., Etiological phytopathology, 843–862
LINK, KARL P., see DICKSON, JAMES G.
Locality, influence of, on sporulation of Phyllosticta solitaria, 503–504
Longevity of sugar-beet mosaic virus in vitro, 469
Losses, due to deterioration of forest products, 13; from cereal diseases, method of estimating, 12; from disease, as indicated by total yield, 983–984
LUDBROOK, W. V., Pathogenicity and environal studies on Verticillium hadromycosis, 117–154
Lupin inoculated with Verticillium dahliae, 117–154
Lycopersicon esculentum, isolation of yellow-mosaic virus from, 965

MACHACEK, J. E., see GREANEY, F. J.; and F. J. GREANEY, Relation of mechanical injury to foot rot of cereals, 22
MACKIE, W. W., Correction, 205
Macrophomina phaseoli, correction in article by W. W. MACKIE, 205
Macrosiphum, ambrosia, vector of bean-mosaic virus, 40; solanifolii, transmission of some viruses affecting spinach, 446–474
MADER, E. O., Effect of pressure and amounts of copper applied in spraying potatoes, 22; Evidence of stimulation of potatoes by Bordeaux mixture, 22; see BLODGETT, F. M.
Maize seedlings, reaction of, to Gibberella saubinetii, 905
MARRITT, J. W., see SANFORD, G. B.
MASSEY, L. M., R. P. WHITE, and A. E. JENKINS, Disease of cultivated sweet violet caused by Sphaceloma, 22–23
MCCALLAN, S. E. A., and FRANK WILCOXON, Form of toxicity surface for copper sulphate and for sulphur in relation to conidia of Sclerotinia americana, 23
MCCLELLAND, C. K., see YOUNG, V. H.
MCCORMACK, R. B., see BLODGETT, F. M.
MCCOWN, MONROE, Weak Bordeaux spray in the control of fire-blight of apple, 729–733

McKenzie, Malcolm A., see Waterman, Alma M.
Mealy bug, evidence of green-spotting strains, 254–256; green spotting, feeding effects on pineapple, 243; hereditary nature of capacity to form green spots, 253–254; hereditary nature of capacity to form nongreen spots, 250–251; nongreen spotting, feeding effects on pineapple, 246; nongreen spotting, relation to pineapple wilt, 234; pineapple, 207
Mechanical injury, relation to foot rot of cereals, 22
Media, humus-extract agar for oospore production in Pythium, 745; influence on sporulation of Phyllosticta solitaria, 505–519
Medicago sativa, mosaic, 562
Medium, Czapek's, used for Helminthosporium sativum, 617
Meier, F. C., J. A. Stevenson, and Vera K. Charles, Spores in the upper air, 23
Melampsora lini, heterothallism in, 4; spermatia of, 487
Melampsoropsis, abietina, on spruce in Minnesota, 615; cassandrae, on spruce needles in Minnesota, 615; ledicola, on spruce needles in Minnesota, 615
Melilotus, alba, mosaic, 562; officinalis, mosaic, 562
Melon, relationship of Fusarium niveum to the formation of tyloses in plants, 33
Mercuric chloride, acidulated, toxicity to sclerotia of Rhizoctonia solani, 275; acidulated, decrease in strength and effectiveness, 275; acidulated, effect on vitality of potato sets, 277; decrease in strength and effectiveness, 275; effect on vitality of potato sets, 277; fungicidal efficiency after addition of potassium iodide to solution of, 28–29; toxicity to sclerotia of Rhizoctonia solani, 273–275; treatment with, for control of gladiolus scab, 804–808
Merko, for control of oat smut, 827
Method of measuring virus concentration, 752
Methods, of estimating losses from plant diseases, 975–984; of inoculating tobacco with virus of typical mosaic, through stomata, 935–938
Microbiology, single-cell isolation in, 357
Microcera sp. in Brazil, 737
Mielke, Jas. L., Comparison of pycnial stage of Cronartium ribicola on Pinus lambertiana and P. monticola, 204–205; Tuberculina maxima in Western North America, 299–305
Mildew, downy, of tobacco on pepper, tomato, and eggplant, 837; powdery, of crape myrtle, 814, 1002–1003; powdery, of crape myrtle, control of, 818
Miles, L. E., Control of gladiolus scab, 802–813
Miller, Edwin C., book review, Principles of plant physiology, by Oran Raber, 1005–1006; see Johnston, C. O.
Minnesota, field observations on needle rusts of spruce in, 613–616
Mitchella repens, intumescences on, 286
Mites, associated with disease, of Datura stramonium, 622; associated with disease, of Nicandra physaloides, 622; associated with disease, of pepper, 622; associated with disease, of Physalis spp., 622; associated with disease, of potato, 622–624; associated with disease, of Solanum antipoviczii hybrids, 622; associated with disease, of S. demissum hybrids, 622; associated with disease, of S. spp., 622; associated with disease, of tomato, 622; of fig. 893; on potato, lime sulphur and sulphur used as preventives, 624
Mix, A. J., Factors affecting the sporulation of Phyllosticta solitaria in artificial culture, 503–524
Monteith, John, Jr., A Pythium disease of turf, 23–24
Moore, Elizabeth Jane, Growth relations in culture of the cotton-root-rot organism, Phymatotrichum omnivorum, 525–537
Morphology of Stagonospora curtisii, 781
Morris, Robert T., A hitherto unreported blight of the Juglandaceae, 407–408
Mosaic, cucumber, inheritability of virus in aphids, 467–468; cucumber, isolation

of virus from naturally infected spinach, 466–467; cucumber, on spinach, resemblance to spinach blight, 461–463; cucumber, on tobacco, 311; cucumber, transmission to spinach, 459–463; diseases, legume, in relation to beans and peas, 562–564; of dahlia, 6; of fig, 887; of fig in California, 7; of flue-cured tobacco, effect on yield and quality, 834; of peach, 17; of tobacco, control of, 831; of tobacco, entrance of the virus through stomata, 934–948; properties of sugar-beet virus, 468–470; rugose, of potato, 319–342; sugar-beet, longevity of virus in vitro, 469; sugar-beet, thermal death point of virus, 470; sugar-beet, tolerance to dilution of virus, 470; sugar-beet, transmission to spinach, 463–465; suspected mosaic of strawberry, 654; tobacco, a source of inoculum for, 839; virus, of tobacco, 749; watermelon, occurrence of, 741–744; yellow, from tobacco, viruses of, characters produced in several inoculated hosts, 970–972; yellow, from tobacco, viruses of considered to be strains of tobacco mosaic, 973; yellow, of tomato, 34; yellow, viruses of, isolation from plants infected with tobacco mosaic, 964–974

Moss, E. G., see Wolf, Frederick A.

Mottle, in potato rugose-mosaic complex, 325, 326, 327, 331–334; tobacco, 325, 326, 329, 330, 331–334

Müller, Albert S., Observations and notes on citrus diseases in Minas Geraes, Brazil, 734–737

Mushrooms, effect of carbon dioxide on, 20

Muskmelon, Fusarium wilt of, in Minnesota, 554–556

Mutation, of Helminthosporium sativum on a no-nitrogen medium, 617–619

Mycelium radicis, atrovirens, true mycorhizal fungus in contrast to, 14; nigrostrigosum, isolated from black mycorhiza of Pinus in Sweden, 14

Mycorhizal fungus, in contrast to Mycelium radicis atrovirens, 14

Myriangium duriaei, in Brazil, 737

Myrtle, crape, powdery mildew of, 814

Mysore, black rot of coffee in, 875

Myzus, fragaefolii, vector of strawberry crinkle, 738–740; persicae, transmission of some viruses affecting spinach, 446–474; persicae, vector of bean-mosaic virus, 40

Narasimham, M. J., Black rot of coffee in Mysore, 875–886; Cytological investigations on spike disease of sandal, Santalum album, 191–202

Narcissus, fire, 773; leaf scorch of, 770; poeticus, inoculation of, with Stagonospora curtisii, 775

Navel-orange scab in Brazil, caused by Sphaceloma fawcettii var. **viscosa**, n. var., 538–545

Neal, D. C., and R. E. Wester, An undescribed sclerotium fungus prevalent in northeast Texas, 24; R. E. Wester, and K. C. Gunn, Fusion of large-cell hyphae of the cotton-root-rot fungus, 676–677; R. E. Wester, and K. C. Gunn, Morphology and life history of the cotton-root-rot fungus in Texas, 24

Nebuka, Allium fistulosum, 292; resistance to pink root, 294–297

Necrosis, in phloem and pericycle, of curly-top-affected beet leaves, localization of, 686

Nectria, associated with root rot of citrus, 735; canker, of basswood, 36; canker, on various trees, 36; coccinea, on beech, 10

Nelson, Ray, and L. C. Cochran, Breeding Fusarium-yellows-resistant celery, 25; and E. E. Down, Influence of pollen and ovule infection in seed transmission of bean mosaic, 25

Nema, the bulb or stem nema, Tylenchus dipsaci, three new hosts, 620; unusual symptoms produced by Aphelenchoides fragariae, 622

Nematospora, coryli, on grapefruit, in Florida, 384; coryli, on orange, in Florida, 384; coryli, on pecans, possibly carried by insects, 1001; coryli, on satsuma oranges, in Florida, 384; coryli, on sweet pepper, in Florida, 384; coryli, on tomato, in Florida, 384; coryli, pecans infected with, 1000–1001; gossypii, first report in United States, 385; gossypii,

on satsuma orange, in Florida, 385; spp., occurrence, in Florida, 384
NEWTON, MARGARET, and A. M. BROWN, Studies on the resistance of wheat and oats to stem rust, 25; T. JOHNSON, and A. M. BROWN, Stripe rust in Canada, 25–26
Nicandra physaloides, affected by mites, 622
NICHOLS, R. A., see BRESSMAN, E. N.
Nicotiana, glutinosa, as test-plant for virus, 752; spp., inoculation of, with viruses of yellow mosaic, 969; spp., isolation of yellow-mosaic viruses from, 965–974; tabacum, anatomy of "Kroepoek-diseased" leaf of, 175–190; tabacum, and sugar beet, comparison of viruses on, 451–458; tabacum, mosaic virus of, 749
Nitrogen, source of, effect on sporulation of Phyllosticta solitaria, 515–517
Nongreen spotting, anatomy of, 247; of pineapple leaves, 246
NORTON, J. B. S., see HEUBERGER, J. W.; and J. W. HEUBERGER, Factors influencing type and sequence of tomato-mosaic leaf abnormalities, 26
Nutrition as a cause of fungus mutation, 617
Nyasaland, new species of Sclerospora from, 587–595

Oat, smut, control of, 825; -smut fungi, inheritance of chlamydospore characteristics in, 16
Oats, resistance to stem rust, 25
Okra, inoculated with Verticillium dahliae, 125
Olea, host of Dothiorella ribus, 929
Olive, development and control of knot, 37
Onion, Allium cepa, resistance to pink root, 292–293; varietal differences, 295, 296
Oospora, citri-aurantii, on citrus in Brazil, 737
Oospore production, Pythium, humus-extract agar for, 745
Oospores, germination of, of Pseudoperonospora humile, 485–487; of **Sclerospora butleri**, n. sp., 587–595
Ophiobolus graminis, ascospore discharge in, and its probable relation to the development of whiteheads in wheat, 721–728; incidence of, in South Australia, 722; longevity of the spores of, 725; spore dissemination of, 725
Orange, Navel, Sphaceloma fawcettii var. **viscosa**, n. var., cause of scab in Brazil, 538–545; Nematospora coryli on, in Florida, 384; satsuma, N. coryli on, in Florida, 384; satsuma, N. gossypii on, in Florida, 385; sour, Sphaceloma fawcettii on, in Brazil, 544; sweet, Sphaceloma on, in Argentina, Brazil, and Paraguay, 544
Orchard grass, sclerotium from, 867
ORTON, C. R., Reaction of tissues from individual plants to soft-rot bacteria, 26; and ALFRED R. STANLEY, Serum-agglutination studies with soft-rot bacteria, 27
Overwintering, of Phytomonas pruni in culture, 795; of P. pruni in leaves, 797; of P. pruni in twigs, 797; of P. pruni on peach, 787
Oxalis, spermatia of corn rus on, 923
Ozonium texanum, n. sp., on cotton stalks and roots in northeast Texas, 24

Panicum, barbinode, host of pineapple mealy bug, 213, 249; grass, host of pineapple mealy bug, 213, 249
Parasitism, discussion of, 855
Pathogenicity, and environal studies, on Verticillium hadromycosis, 117–154; of **Diaporthe crotalariae**, n. sp., 601–603; of Rhizoctonia bataticola, at high temperatures, 949–963; of V. albo-atrum, 125–128; of V. dahliae, 125–128; of Wojnowicia graminis, 1001–1002
Pathologic changes in anatomy of leaves of sugar beet affected by curly top, 679–712
Pathology, of Chionodoxa luciliae infested by Tylenchus dipsaci, 620; of Chrysanthemum hortorum infested by Aphelenchoides fragariae, 622; of Colchicum speciosum album infested by T. dipsaci, 620; of Digitalis purpurea infested by T. dipsaci, 620; of potato disease due to mites, 622–624; of tobacco black shank, 605–612; plant, contribution to land-utilization program, 404–406
PAXTON, G. E., Consistent mutation of

Helminthosporium sativum on a no-nitrogen medium, 617–619
Pea, relation of different legume mosaics to, 564; sweet, relation of different legume mosaics to, 564; wilt, seed-borne bacteria main cause, 14
Peach, Armillaria mellea on, 28; Bacterium tumefaciens on, 28; brown rot, distribution of losses from, in the United States, 978; brown rot, in cars, in the United States, 980; brown rot, losses from, in the United States, 979; description of spring cankers of Phytomonas pruni on, 787; description of summer cankers of P. pruni on, 787; Heterodera radicicola on, 28; mosaic, 17; overwintering of P. pruni on, 787; Polyporus curtisii cause of heart rot, 28
Peanut, Arachis hypogaea, leaf spot, 627–640
Peas, influence of root rot on, 16–17; relation of soil fertility to incidence of Aphanomyces root rot of, 36; studies of varieties resistant and susceptible to Fusarium wilt, 36
Pecans, Nematospora coryli, on kernels, 1000–1001
Pellicularia koleroga, synonym of Corticium koleroga, 876
PELTIER, GEORGE L., Physiologic forms of wheat stem rust in Kansas and Nebraska, 343–356
Penicillium, digitatum, on citrus in Brazil, 737; effect of vitamins on growth of, 929; italicum, on citrus in Brazil, 737
Pennisetum typhoideum, Sclerospora graminicola on, in Nyasaland, 587, 593
Pentathionates, physical properties of, 159–161; preparation of, 158–159; quantitative analysis of, 161–163
Pentathionic acid, and its salts, 155–174; preparation and physical properties of, 163; toxicity of, 168–171
Pepper (Capsicum sp.), affected by mites, 622; downy mildew of tobacco on, 837; inoculated with Verticillium albo-atrum, 124, 127; inoculated with V. dahliae, 124, 125, 127; Nematospora coryli on, in Florida, 384
Periderm, formation in intumescences, 289
Peronospora, effusa, infection of seed clusters of spinach by, 7; frequency distribution of spore size of, 838; hyoscyami, on tobacco, pepper, tomato, and eggplant, 837; schleideni, resistance in Allium fistulosum to, 109–110
Persea, americana, Dothiorella ribis from, 929; host of D. ribis, 929
PERSON, L. H., A bacterial blight of the broad bean in Louisiana, 27
Phaseolus, limensis, resistance to Rhizoctonia bataticola, 954; lunatus macrocarpus Benth., scab disease, 622; lunatus, resistance to Rhizoctonia bataticola, 953; vulgaris, as test plant for virus, 752; vulgaris, environmental factors in relation to transit diseases of, 411–445; vulgaris, mosaic, 562; vulgaris, Rhizoctonia bataticola causing seedling stem blight of, 949–963
Pheidole megacephala, ant, attendant of pineapple mealy bug, 213
Phenology, new journal of, 316
Phoma terrestris, cause of pink root of Allium, 290; resistance in A. fistulosum to, 109–110; resistance of cultivated species of Allium to, 290–298
Phomopsis, callistephi, cause of stem canker of China aster, 20; **crotalariae**, n. f. nom., imperfect stage of **Diaporthe crotalariae**, n. sp.,600, 604; on citrus in Brazil, 735
Phyllactinia corylea, on crape myrtle, comparison with Erysiphe lagerstroemiae, 1002–1003; on crape myrtle in Alabama, 1003
Phyllosticta, causing leaf spot of citrus in Brazil, 736; effect of vitamins on growth of, 929; solitaria, autolysis in culture, 522; solitaria, cardinal temperatures, 521; solitaria, factors affecting sporulation in artificial culture, 503–524; solitaria, new culture medium for, 514; solitaria, nonsporulating strains, 522; solitaria, saltation in culture, 522; solitaria, sporulating strains, 521–522; solitaria, sporulation of, effect of carbon source on, 518–519; solitaria, sporulation of, effect of hydrogen-ion concentration on, 519; solitaria, sporulation of, effect of nitrogen source on, 515–517; solitaria,

sporulation of, effect of temperature on, 519–521; solitaria, sporulation of, influence of food quality on, 511–515; solitaria, sporulation of, influence of food quantity on, 506–515; solitaria, viability of spores from culture, 523

Phymatotrichum omnivorum, growth, on barley-root-decoction medium, 533–534, 536; growth, on corn-root-decoction medium, 529–533, 535; growth, on cotton-root-decoction medium, 529–533, 535; growth, on potato dextrose, 529–533, 535; growth, on Turk's cap-root-decoction medium, 531–533, 535; growth, on wheat-root-decoction medium, 533–534, 536; growth relations in culture, 525–537; morphology and life history in Texas, 24; resistance to, relation to growth-inhibiting substances, 536; (Shear) Duggar, fusion of hyphae, 676; toxicity of barley-root decoction to, 533–534, 536; toxicity, of corn-root decoction to, 529–533, 535; toxicity, of Turk's cap-root decoction to, 531–533, 535; toxicity, of wheat-root decoction to, 533–534, 536

Physalis, angulata, inoculation of, with viruses of yellow mosaic from tobacco, 969; spp., affected by mites, 622

Physalospora, rhodina, hosts of, 70–72; zeicola, hosts of, 70–72; zeicola, on corn, 63–72; zeicola, relation to Diplodia frumentia, on corn, 63–68; zeicola, taxonomic and host relationships, 63–72

Physiologic, forms, distinguished in Tilletia tritici by differences in size and shape of smut balls, 16; forms, of Bremia lactucae on lettuce, 18–19; forms, of Colletotrichum falcatum, 557–559; forms, of Puccinia graminis tritici in relation to stem-rust epidemiology in 1932, 34; forms, of stripe rust in Canada, 25; forms, of Tilletia tritici, wheat varieties resistant to, 932; forms, of wheat stem rust, 343–356; specialization, of Tilletia tritici on emmer, 483–485

Phytomonas, pruni, die-back of twigs caused by, 790; pruni, infection of primary foliage by, 789; pruni, isolations of, from overwintering leaves, 797; pruni, overwintering of, in culture, 795; pruni, overwintering of, in twigs, 797; pruni, overwintering on peach, 787; pruni, resistance of basal portions of twigs to, 798; tumefaciens, causing crown gall of Cupressus arizonica, 97–101

Phytopathological and botanical research methods, book review, 1004–1005

Phytopathology, etiological, 843

Phytophthora, associated with damping-off of citrus, 735; associated with gum diseases of citrus, 734; citrophthora, on Citrus in Florida and Louisiana, 667–669; hibernalis, on Citrus, in California, 667–668; infestans, culture of, 713–720; infestans, sporulation, temperature range of, 715; nicotianae, acid production by, 610; nicotianae, cause of tobacco black shank, 605–612; nicotianae, growth in artificial cultures, 606, 608; nicotianae, influence of pH concentration on wilting, 610, 611; nicotianae, nature of toxin, 609; nicotianae, toxin in diseased plants, 608, 609; nicotianae, toxin production in cultures, 608, 609; nicotianae, utilization of carbohydrates, 611; parasitica, 734; parasitica, on Citrus, in Florida, 667, 668

Picea, excelsa, susceptibility to needle rusts in Minnesota, 513; glauca albertiana, susceptibility to needle rusts in Minnesota, 613; glauca, susceptibility to needle rusts in Minnesota, 613; mariana, susceptibility to needle rust in Minnesota, 613; pungens, susceptibility to needle rust in Minnesota, 613; spp., Cytospora, canker of, 11; spp., susceptibility to needle rusts in Minnesota, 613

PIERSTORFF, A. L., and HOWARD LAMB, Dissemination of fire-blight bacteria by honeybees, 27

Pine, white, blister rust, authority for name or organism, 559–561

Pineapple, leaf spot, 7; mealy bug, green spotting and nongreen spotting, relation to pineapple wilt, 234; mealy bug, hosts, 213; mealy bug, Pseudococcus brevipes, 207; mealy bug, relation of ants to, 213–217; mealy bug, relation to pineapple wilt, 213–235; mealy bug, relation to spotting of pineapple leaves, 243; mealy

bug, symbionts, possible relation to green spotting, 256–258; mealy bug, toxicity of, in relation to original hosts, 228–234; mealy bug, types of spotting from, 243; wilt of, 207; wilt, description of, 208–211; wilt, edge wilt, synonym of, 208; wilt, possible nature of injury, 239; wilt, recovery from, 235; wilt, relation of age of plants to susceptibility, 226; wilt, relation of green spotting to, 234; wilt, relation of pineapple mealy bug to, 213–235

Pink root, of Allium, 290; resistance of cultivated species of Allium to, 290–298; susceptibility of various species to, 295

Pinus, lambertiana, pycnial stage of Cronartium ribicola on, 204–205; monticola, pycnial stage of C. ribicola on, 204–205; monticola, resistance of current season's shoots of, to infection by C. ribicola, 917; spp., mycorhiza, 14; strobus, field inoculations with sporidia of C. ribicola in Minnesota, 105–107

PIRONE, P. P., Combating spinach damping-off by seed treatment, 28

Pisum sativum, mosaic, 562

Plant, diseases, in the United States, losses from, 975–984; pathology, contribution to land-utilization program, 404–406

Podonectria sp. in Brazil, 737

Pokkah-bong disease, hosts other than sugarcane, 672

Polymyxa graminis, life history, morphology, and cytology of, 20–21

Polyporus curtisii on peach trees in the Carolinas, 28

Polystictus versicolor, growth as affected by temperature, 73–81

Polythionic acids, analyses, 163–165

POOLE, R. F., Knob and elevated vein formations on sweet-potato roots, 91–96; Prevalence and relations of four soil-borne parasites on the peach trees in the Sand Hills of the Carolinas, 28

Populus grandidentata, Nectria canker of, 36.

PORTER, D. R., and H. A. JONES, Resistance of some of the cultivated species of Allium to pink root (Phoma terrestris), 290–298

Portulaca oleraceus, host of pineapple mealy bug, 249

Potassium iodide, use in mercuric chloride solution, 28–29

Potato, a new necrotic virus disease of, 32; Botrytis cinerea, inoculation with, 995–997; disease associated with mites, in New York, 622–624; effect of pressure and amounts of copper applied in spraying, 22; influence of spray on temperature of, 912; late blight, control of, 29; latent mosaic, resistance to, 32; leaf hopper, nature of injury on forage legumes, 19; Rhizoctonia, mercuric chloride and potassium iodide solution for control of, 28–29; rugose mosaic, nature of, 319–342; scab, environment in relation to, 9; stimulation by Bordeaux mixture, 22; susceptibility of, to Verticillium alboatrum, 125; susceptibility of, to V. dahliae, 125; tuber rot, caused by Botrytis cinerea, 11, 993–999; Verticillium hadromycosis, 143–146

Powdery mildew of crape myrtle, 814

Preparation and properties of pentathionic acid and its salts, its toxicity to fungi, bacteria, and insects, 155–174

PRICE, W. C., The thermal death rate of tobacco-mosaic virus, 749–769

Principles of plant physiology, book review, 1005–1006

PRIODE, C. N., Cuban streak, 674–676; Two hosts of pokkah-bong disease other than sugarcane, 672–674

Promycelia of Puccinia malvacearum, cytology of, 572–586

Properties and preparation of pentathionic acid and its salts, its toxicity to fungi, bacteria, and insects, 155–174

Prunus, serotina, Nectria canker of, 36; supp., bacterial canker in California 36–37; spp., hosts of Dothiorella ribis, 929

Pseudococcus brevipes, 207; relation to pineapple wilt, 213; to green spotting, 259; to nongreen or chlorotic spotting, 259

Pseudoperonospora humuli, germination of oospores of, 485–487

Psidium, host of Dothiorella ribis, 929

Psorosis, a virus disease of Citrus, 930

Psyllid yellows, differentiated from curly top, 930; insect toxin theory of etiology of, 930; of tomato, 930
Puccinia, coronata, heterothallism in, 4; glumarum tritici, in Canada, 25; graminis, heterothallism in, 4; graminis tritici, physiologic forms in relation to stem-rust epidemiology in 1932, 34; graminis tritici, physiologic forms of, 343; graminis tritici, resistant wheats from Russia, 8; malvacearum, behavior of nucleole in nuclear division, 582; malvacearum, chromosome number of, 581; malvacearum, cytology of teliospores, promycelia and sporidia, 572–586; malvacearum, existence of centrosome in nucleus, 581; sorghi, spermatia of, 923; triticina, resistant wheats from Russia, 8
Pure-culture technique for quantitative studies of plant growth in association with microorganisms, 14
Pycnidia, of **Diaporthe crotalariae**, n. sp., 596, 600–602; of Stagnospora curtisii, 778, 783
Pyracantha, host of Dothiorella ribis, 929
Pythium, arrhenomanes, 745; butleri, cause of a disease of turf, 23–24; butleri, on snap beans in transit, 428; oospore production, humus-extract agar for, 745

Quality, effect of tobacco mosaic on, 834
QUANJER, H. M., A complex virosis of tobacco, 28
Quercus, host of Dothiorella ribis, 929
Quick wilt, pineapple, 211
Quince-rust, caused by Gymnosporangium germinale, 546–553; etiology of, 549–552

RABER, ORAN, Principles of plant physiology, book review, 1005–1006
RALEIGH, W. P., Fungicidal efficiency of a solution of mercuric chloride and potassium iodide, 28–29; A homemade colloidal copper spray, 29; see SCHULTZ, E. S.
RANDS, R. D., and ERNEST DOPP, Humus-extract agar favorable for oospore production in Pythium, 745
RANKIN, W. H., see FELT, E. P.
Raspberry, bacterial twig and blossom blight of, 21; inoculated with Verticillium dahliae, 125
RAWLINS, THOMAS ELLSWORTH, Phytopathological and botanical research methods, book review, 1004–1005; see TAKAHASHI, WILLIAM N.
Red cedar (see also cedar, red), Juniperus virginiana
Red-top grass, host of pineapple mealy bug, 213, 249
REDDICK, D., A potato disease, 622–624
Rehmiellopsis bohemica, attacking Abies concolor, 108–109
REMSBERG, RUTH, and C. W. HUNGERFORD, Certain Sclerotium diseases of grains and grasses, 863–874; see HUNGERFORD, C. W.
Report of the twenty-fourth annual meeting of The American Phytopathological Society, 489–500
Resistance, in Allium fistulosum to Urocystis cepulae, Phoma terrestris, and Peronospora schleideni, 109–110; of basal portions of twigs to Phytomonas pruni, 798; of corn to seedling blight, nature of, 9; of cultivated species of Allium to pink root, 290–298; of potato to latent mosaic, 32; of rosaceous plants to fire blight, 32–33; to Fusarium wilt in peas, 36; to Phymatotrichum omnivorum, relation to growth-inhibiting substances, 536; to rust in beans, 38
Rhizoctonia, bataticola, causing seedling stem blight of field beans at high temperatures, 949–963; bataticola, correction in article by W. W. MACKIE, 205; bataticola, cultures of, 957–961; bataticola, on bean, distribution of, in California, 949–963; bataticola, on bean, economic importance of, in California, 949–963; solani, 271; solani, effect of acidulatid mercuric chloride on sclerotia, 275; solani, effect of formaldehyde on sclerotia, 272; solani, effect of mercuric chloride on sclerotia, 273; solani, effect of temperature on brown patch caused by, 8; solani, on snap beans in transit, 428–431
Rhizopus, nigricans, on snap beans in

transit, 431–436; soft rot on snap beans in transit, 431–436; tritici, on snap beans in transit, 431–436
Rhus typhina, inoculated with Verticillium dahliae, 125; Nectria canker of, 36
Rice, genetic connection between Sclerotium oryzae, Helminthosporium sigmoideum, and Leptosphaeria salvinii on, 35
RICHARDS, B. L., see BLOOD, H. L.
RICHARDS, L. A., see BLOOD, H. L.; and H. L. BLOOD, Some improvements in auto-irrigator apparatus, 928–929
RIKER, A. J., and E. M. HILDEBRAND, Seasonal development of hairy root, crown gall, and wound overgrowth on apple trees in the nursery, 29; G. W. KEITT, E. M. HILDEBRAND, and W. M. BANFIELD, Control of hairy root, crown gall, and other malformations at the unions of piece-root-grafted apple trees, 30
RIKER, REGINA, see JONES, L. R.; and L. R. JONES, Fusarium strains in relation to wilt in China aster, 29–30
Ring, blotch, of citrus, possible virus origin of, 930; spot, in potato rugose-mosaic complex, 327–331, 332, 333, 334; -spot-like virus disease of red clover, 746–747; spot, of dahlia, 6; spot, tobacco, transmission to spinach, 466; spot, yellow, of dahlia, 6
Robinia, pseudoacacia, brooming disease transmitted by grafts, 13; pseudoacacia, transmissibility of brooming disease of, 83–90
RODENHISER, H. A., The genetics of Sphacelotheca sorghi and S. cruenta, 30; and B. F. BARNES, Pathogenicity of certain hybrids of covered and loose smuts of sorghum, 30–31
Roguing in the control of tobacco mosaic, 831
ROLLINS, R. C., see SOLHEIM, W. G.
Rosa, host of Dothiorella ribis, 929
Rose, inoculated with Verticillium dahlias, 125
ROSEN, H. R., Influence of spray applications on air temperatures surrounding sprayed potato plants, 912–916
Root-knot nematode, control of, in greenhouses with carbon disulphide emulsion, 13
Root-rot, fungus, of cotton, fusion of hyphae, 676; influence on peas, 16–17
Rubus, host of Dothiorella ribis, 929
RUDOLPH, B. A., Report of the seventeenth annual meeting of the Pacific Division of The American Phytopathological Society, 928
RUNNELS, H. A., see J. D. WILSON
Rust, apple, caused by Gymnosporangium juniperi-virginianae, 546; hawthorn, caused by G. globosum, 546; quince, caused by G. germinale, 546–553; spermatia of, on flax, 487
Rusts, centrosome in nucleus, 581; chromosome number of, 580, 581; cytology of Puccinia malvacearum, 572–586; nucleole and its behavior in nuclear division, 582; of spruce needles in Minnesota, field observations, 613–616
RUTTLE-NEBEL, MABEL L., Comparative studies of field collections of Ustilago hordei and U. nuda, 31

Sahuaro (Carnegiea gigantea), crown gall on, 21
Salix, host of Dothiorella ribis, 929
Saltant of Helminthiosporium sativum produced by ultra-violet radiation, 379
Saltation, in Helminthosporium sativum, produced by ultra-violet radiation, 12; of Phyllosticta solitaria in culture, 522
SAMUEL, GEOFFREY, and S. D. GARRETT, Ascospore discharge in Ophiobolus graminis, and its probable relation to the development of whiteheads in wheat, 721–728; J. G. BALD, and C. M. EARDLEY, "Big bud," a virus disease of the tomato, 641–653
Sandal, cytological investigations on spike disease of, 191–202
SANDERS, GEORGE E., Notes on Bordeaux mixture, 31–32
SANFORD, G. B., and J. W. MARRITT, The toxicity of formaldehyde and mercuric chloride solutions on various sizes of sclerotia of Rhizoctonia solani, 271–280
Santalum album, cytological investigations on spike disease of, 191–202
Scab, citrus, distribution of, 481; gladiolus, control of, 802; gladiolus, control of, by spraying and dusting, 812; Navel orange,

in Brazil, 538–545; of Lima bean, geographic range, 662; of Lima bean, life history, 665; sweet orange, in Argentina, distinct from common citrus scab, 479; use of term as applied to diseases caused by Sphaceloma and Gloeosporium, 389–395

Scale, Mediterranean fig, in California, 894

Scaly bark of citrus, 930

SCHULTZ, E. S., and W. P. RALEIGH, A new necrotic virus disease of potatoes, 32; Resistance of potato to latent mosaic, 32

SCHUSTER, C. E., A disorder of Persian (English) walnuts grafted on black-walnut stocks, resulting in girdling, 408–409

Sclerospora, **butleri**, characteristics of fungus, 589–591; **butleri**, economic importance of host, 594; **butleri**, effect of fungus on host, 589; **butleri**, identity of fungus, 590–593; **butleri**, new species from Nyasaland, 587–595; **butleri**, occurrence, 587; **butleri**, technical description, 592, 593; **graminicola**, on Penisetum typhoideum in Nyasaland, 587, 593

Sclerotia, from grains and grasses, 867; of Corticium koleroga, 881; Rhizoctonia solani, relation of size to killing by fungicides, 273–274; R. solani, relation of time of action to killing by fungicides, 273–274; R. solani, size in nature, 277

Sclerotinia, americana, form of toxicity surface for copper sulphate and for sulphur in relation to conidia of, 23; **gladioli**, n. comb., perfect stage of Sclerotium gladioli, 9; libertiana, causing stem canker of hollyhock, 39; sclerotiorum, on snap beans in transit, 425–428

Sclerotium, diseases, of grains and grasses, 17, 863; gladioli, hosts of, 9; **gladioli**, n. comb., perfect stage of, 9; oryzae, ascigerous stages is Leptosphaeria salvinii, 35; rhizodes, snow mold and brown patch caused by, 8; rolfsii, on snap beans in transit, 431

Scorch, leaf, of narcissus, 770

SEARS, S. S., see SOLHEIM, W. G.

Seed, transmission, of bean mosaic, 25; transmission, of legume-mosaic viruses, 563; transmission, of Sorosporium reilianum, 398; transmission, of virus causing variegation of Abutilon, 20; treatment, combating spinach damping off by, 28; treatment, in relation to common potato scab, 9

Seedling stem blight of field beans caused by Rhizoctonia bataticola at high temperatures, 949–963

Seedlings, use of formaldehyde dust to control damping off, 38

Semesan, treatment for control of gladiolus scab, 804, 805, 810

Septogloeum arachidis Rac., synonym of Cercospora personata (B. & C.) E. & E., 637

Serological reaction, as means of determining concentration of tobacco-mosaic virus, 4

Serum agglutination, with soft-rot bacteria, 27

Shallots, Allium ascalonicum, resistance to pink root, 294

SHANDS, R. G., see DICKSON, JAMES G.

SHAPOVALOV, M., Dieback form of tomato streak, 928

SHAW, LUTHER, Resistance of rosaceous plants to fire blight, 32–33

SHEAR, C. L., NEIL E. STEVENS, and HENRY F. BAIN, Fungous diseases of the cultivated cranberry, review of, 314–315

SHEAR, ELMER V., and J. S. COOLEY, Relation of growth cycle and nutrition to perennial apple-canker infection, 33

Shipment, environmental factors in relation to snap-bean diseases occurring in shipment, 411–445

Single-cell isolation in microbiology, 357

Sisal, host of pineapple mealy bug, 213, 249

SLEETH, BAILEY, Relationship of Fusarium niveum to the formation of tyloses in melon plants, 33

SMITH, C. O., Inoculation with Dothiorella ribis, 929

Smut, head, of corn, in western Oregon, 396–403; oat, control of, 825; physiologic specialization of Tilletia tritici on emmer, 483–485

Snap bean, environmental factors in rela-

tion to diseases occurring in shipment, 411–445
Snow mold, caused by Sclerotium rhizodes, 8; of grains and grasses, 864
Snow, rot, of grains and grasses, 864; scald, of grains and grasses, 864
Soft-rot bacteria, reaction of tissues from individual plants to, 26; serum agglutination with, 27
Soil, contamination, with Sorosporium reilianum, 400–403; fertility, relation to incidence of Aphanomyces root rot of peas, 36; infestation, by chlamydospores of Tilletia levis in Montana, 39; moisture, influence on infection of wheat by Urocystis tritici, 10–11; rot, on snap beans in transit, 428–431; temperature, influence of infection of wheat by Urocystis tritici, 10–11
Solanum, antipoviczii, hybrids, affected by mites, 622; demissum, hybrids, affected by mites, 622; melongena, inoculation of, with viruses of yellow mosaic from tobacco, 969; spp., affected by mites, 622; tuberosum, Botrytis cinerea on, 933–999
Solenopsis geminata, var. rufa, ant, attendant of pineapple mealy bug, 213
SOLHEIM, W. G., S. S. SEARS, and R. C. ROLLINS, Effect of vitamins on growth of fungi in pure culture, 929–930
Sorghum, a host of pokkah-bong, 672; observations on Tolyposporium filiferum, cause of "long smut" of, 985–992
Sorosporium reilianum on corn in western Oregon, 396–403
Source of culture, influence of, on sporulation of Phyllosticta solitaria, 504
Southern blight on snap beans in transit, 431
SPAULDING, PERLEY, Dietrich preferable authority for Cronartium ribicola, 203–204
Speck, bacterial, of tomato, 897
Spermatia, of flax rust, Melampsora lini, 487; of Puccinia sorghi, 923
Sphaceloma, cruenta, genetics of, 30; cruenta, pathogenicity of certain hybrids with S. sorghi, 30–31; disease of cultivated sweet violet caused by, 22–23; diseases caused by, "anthracnose" and "scab" applied to, 389–395; fawcettii Jenkins, records of, from herbarium specimens of the genus Citrus in England and the United States, 475–482; fawcettii, on sour orange, in Brazil, 544; fawcettii, var. **viscosa**, n. var., cultural studies, 539–542; fawcettii, var. **viscosa**, n. var., description, 543; fawcettii, var. **viscosa**, n. var., on Navel orange in Brazil, 538–545; on sweet orange, in Argentine, Brazil, and Paraguay, 544; sorghi, genetics of, 30; sorghi, pathogenicity of certain hybrids with S. cruenta, 30–31
Spike disease of sandal, cytological investigations on, 191–202
Spinach, blight, resemblance to cucumber mosaic on spinach, 461–463; comparison of viruses on, 459–466; infection of seed clusters by Peronospora effusa, 7; natural infection with cucumber mosaic, 466; seed treatment for damping off of, 28; some viruses affecting, and certain aspects of insect transmission, 446–474; transmission of cucumber mosaic to, 459–463; transmission of sugar-beet mosaic to, 463–465; transmission of tobacco ring spot to, 465–466
Spinacia oleracea, some viruses affecting, and certain aspects of insect transmission, 446–474
Spore size of Peronospora, frequency distribution of, 838
Spores, in upper air, 23; of Tolyposporium filiferum, germination of, 986–989; of Wojnowicia graminis, measurements of, 1002
Sporidia of Puccinia malvacearum, cytology of, 572–586
Sporulation of Phyllosticta solitaria in artificial culture, factors affecting, 503–524
Spot necrosis, tobacco, 323, 326, 330
Spotted wilt, similarity of tomato streak to, 928
SPRAGUE, RODERICK, Association of Cercosporella foot rot with a specific ecological area, 33–34
Spray, applications, influence of, on temperatures of sprayed plants, 912; materials, influence on transpiration, 37
Spraying for control of gladiolus scab, 812
Spruce, field observations on needle rusts in Minnesota, 613–616
Stagonospora curtisii, illustration of pyc-

nidia of, 778, 783; life history of, 775; morphology of, 781; on narcissus, 773; parasitism of, 774; spores of, 779

STAKMAN, E. C., see HAYES, H. K.; LEE HINES, HARRY G. UKKELBERG, and WALLACE BUTLER, Distribution of physiologic forms of Puccinia graminis tritici in relation to stem-rust epidemiology in 1932, 34

STANLEY, ALFRED R., see ORTON, C. R.

STEINER, G., and EDNA M. BUHRER, The bulbous irises as hosts of Tylenchus dipsaci, the bulb or stem nema, 103–105; Three new hosts for Tylenchus dipsaci, the bulb or stem nema, 620; Unusual disease symptoms produced by Aphelenchoides fragariae, 622

Stem rust, of wheat, physiologic forms, in relation to epidemiology in 1932, 34; resistance of oats to, 25; resistance of wheat to, 35

STEVENS, NEIL E., How plant pathology can contribute to a land-utilization program, 404–406; see SHEAR, C. L.; Some significant estimates of losses from plant diseases in the United States, 975–984

STEVENSON, J. A., see MEIER, F. C.

Stilbum cinnabarinum, cause of a new fig disease in Louisiana, 35

Stimulation of potatoes by Bordeaux mixture, 22

Stomata, infection through, with the virus of typical tobacco mosaic, 934–948

Storage, development of apple scab in, 5; rots, sweet potato, losses from, in the United States, 981–983

STOVER, W. G., and M. T. VERMILLION, Some experiments with a yellow mosaic of tomato, 34

Strains, nonsporulating, of Phyllosticta solitaria, 521–522; pathogenic, of Gymnosporangium germinale, 549; sporulating, of P. solitaria, 521–522

Strawberry, crinkle disease, transmission of, 738–740; suspected mosaic, 654–661; wilt, caused by Colletotrichum fragariae, 6

Streak, tomato, dieback form of, 928

Stripe rust in Canada, 25–26

Sugar, beet, and tobacco, comparison of viruses on, 451–458; beet, curly top, pathologic changes in anatomy of leaves of, 679, 712; beet, mosaic, longevity of virus in vitro, 469; beet, mosaic, properties of virus, 468–470; beet, mosaic, thermal death point of virus, 470; beet, mosaic, tolerance to dilution of virus, 470; beet, mosaic, transmission to spinach, 463–465; cane, Cuban streak disease of, 674; cane, red rot, physiologic forms of organism, 557–559

Sulphur, colloidal, 165–168; colloidal, physical properties of, 167; colloidal, preparation of, 166–167; colloidal, toxicity of, 168–173; form of toxicity surface in relation to conidia of Sclerotinia americana, 23

Sulphuric acid, effect on soil pH, 18

Sunlight, effect of, on Sclerotium, 870

Susceptibility, of Allium species to pink root, 290–298; of apple varieties to quince rust, 547; of Picea spp. to needle rusts, 613; of red cedar to Gymnosporangium spp., 546–547; of various hosts to Verticillium hadromycosis, 124–125; varietal, of fig, to mosaic, 889

Suspected mosaic, genetic origin of, 660; noninfectious chlorosis of, 654; of strawberry, occurrence, 654

Sweet, orange, scab, in Argentina, distinct from common citrus scab, 479; pea, inoculated with Verticillium dahliae, 117–154; potato, knob and elevated vein formations on roots of, 91;96; potato, storage rots, losses from, in the United States, 981

TAKAHASHI, WILLIAM N., and THOMAS ELLSWORTH RAWLINS, Evidence regarding the shape of tobacco-mosaic-virus particles, 34–35

Take-all, incidence of, in South Australia, 722

TAPKE, V. F., Comparative studies of two loose smuts of barley (title only), 35

Tarsonemus sp., mite associated with potato disease, in New York, 624

Technique, single-cell isolation, 357

Teliospores of Puccinia malvacearum, cytology of, 572–586

Temperature, as influenced by spray applications, 912; effect of, on cultures of Rhizoctonia bataticola, 961; effect of, on Sclerotium, 870; effect of, on tobacco-mosaic virus, 749; effect on brown patch of turf, 8; influence on sporulation of Phyllosticta solitaria, 519–521; relations, of Verticillium hadromycosis, 146–149

Temperatures, high, effect of, on seedling stem blight of field beans caused by Rhizoctonia bataticola, 949–963

THAXTER, ROLAND (1858–1932), his influence on plant pathology, 565–571; memorial, 502

Thermal, death point, of Heterodera radicicola, 260; death point, of H. radicicola, method of determination, 261; death point, of H. radicicola, relation to time, 262; death point, of sugar-beet-mosaic virus, 470; death rate, of tobacco-mosaic virus, 749

Thielaviopsis paradoxa, not the cause of pineapple leaf spot in Puerto Rico, 7

THOMAS, H. E., The quince-rust disease caused by Gymnosporangium germinale, 546–553

THORNBERRY, H. H., and H. W. ANDERSON, Overwintering of Phytomonas pruni on peach, 787–801

Thurberia thespesioides, intumescences on, 287

TILFORD, P. E., see WILSON, J. D.

Tilia americana, Nectria canker of, 36

Tilletia, levis, soil infestation by chlamydospores of, 39; tritici, differences in size and shape of smut balls of two physiologic forms of, 16; tritici, physiologic specialization, on emmer, 483–485; tritici, resistance of wheat varieties to, 932

Time of isolation, influence on sporulation of Phyllosticta solitaria, 504

Timothy, sclerotium from, 867

TIMS, E. C., Stilbum cinnabarinum, the cause of a new fig disease in Louisiana, 35

Tipburn, relation of Bordeaux spray to, 916

Tobacco, a complex virosis of, 28; and sugar beet, comparison of viruses on, 451–458; cucumber mosaic on, in Wisconsin, 311; downy mildew of, on pepper, tomato, and eggplant, 837; effect of mosaic on yield and quality of, 834; flue-cured, a source of inoculum for mosaic in, 839; Kroepoek, a virus disease of, 175; manufactured, a source of inoculum for mosaic in flue-cured tobacco, 839; mosaic, acquired immunity in, 5; mosaic, control of, 831; mosaic, further purification of virus of, 35; mosaic, isolation of yellow-mosaic viruses from, 964–974; -mosaic virus, evidence regarding shape of particles of, 34–35; -mosaic virus, serological reaction as means of determining the concentration of, 4; -mosaic virus, thermal death rate of, 749; mottle, 325, 326, 329, 330, 331–334; pathology of tobacco black shank on, 605–612; ring spot, intracellular bodies in, 38; ring spot, transmission to spinach, 465–466; ring-spot virus, resistance to aging increased by use of carbolic acid, 14–15; spot necrosis, 323, 326, 330; spot necrosis, acquired immunity in, 5; typical mosaic, entrance of the virus through stomata, 934–948; veinbanding, 323–325, 326, 334; viruses, differentiation of green- and yellow-mosaic, 10

Tolerance to dilution of sugar-beet-mosaic virus, 470

Tolysporium filiferum, cause of "long smut" of sorghum, observations on, 985–992; cultural characters of, 990–992

Tomato, bacterial speck of, 897; big-bud virus disease, 641; blossom-end rot, increased by Bordeaux mixture, 37; curly top of, 929; dieback form of streak of, 928; downy mildew of tobacco on, 837; inoculated with Verticillium albo-atrum, 124–128; inoculated with V. dahliae, 124, 125, 126, 127, 128; (Lycopersicum), affected by mites, 622; mosaic, factors influencing type and sequence of leaf abnormalities, 26; mosaic, water loss in, 15; Nematospora coryli on, in Florida, 384; psyllid yellows of, 930; streak, 34; V. hadromycosis, 138–143; yellow mosaic, some experiments with, 34

Toxicity, of colloidal sulphur, 168–173; of pentathionic acid, 168–171; surface,

form of, for copper sulphate and sulphur in relation to conidia of Sclerotinia americana, 23; to Phymatotrichum omnivorum, of various root decoctions, 529–535, 536
Toxin produced by Phytophthora nicotianae, in diseased plants, 608, 609; in culture media, 606, 608; nature of, 609
Transit diseases of snap beans, environmental factors in relation to, 411–445
Transmission, of cranberry false blossom, by natural graft, 670; of the crinkle disease of strawberry, 738–740
Transpiration, influence of spray materials on, 37
Treatment of gladiolus corms for scab, 803
Trees, cork-borer method for inoculating, 487–488
Tricholaena rosea, host of pineapple mealy bug, 213, 249
Trifolium, pratense, mosaic, 562; pratense, ring-spot-like virus disease of, 746–747; repens, mosaic, 562
Triticum, persicum var. stramineum, resistance to disease, 8–9; timophavii, resistance to disease, 8–9
TRURAN, W. E., see H. F. BERGMAN
Tuberculariа coccicola, 737
Tuberculina maxima, behavior in British Columbia, 301–302; hosts of, 299; on Cronartium comptoniae in British Columbia, 299; on C. ribicola in British Columbia, 299; range in British Columbia, 300; relation to control of white-pine blister rust, 302–304
TULLIS, E. C., Leptosphaeria salvinii, the ascigerous stage of Sclerotium oryzae, 35
Turf, a Pythium disease of, 23–24; brown patch, effect of temperature on, 8
Turk's cap, root decoction, toxicity to Phymatotrichum omnivorum, 531–533, 535
Tylenchus dipsaci, bulb or stem nema, new hosts for, 620; bulbous irises as hosts of, 103–105; on Chionodoxa luciliae, from Holland, 620; on Colchicum speciosum album, from Holland, 620; on Digitalis purpurea, in Connecticut, 620
Tyloses, relationship of Fusarium niveum to formation in melon plants, 33
Typhula graminum, similarity of Sclerotium to, 871
UKKELBERG, HARRY G., see STAKMAN, F. C.
Ulmus, host of Dothiorella ribis, 929
Ultra-violet, light, effect of, on Sclerotium, 870; radiation, of Helminthosporium sativum, producing saltant, 379; radiation, production of white fertile saltant of H. sativum by means of, 12
Uncinula australiana, 1003
Uredo ribicola, 841
Urocystis, cepulae, resistance in Allium fistulosum to, 109–110; tritici, influence of soil moisture and soil temperature on infection of wheat by, 10–11
Ustilago, avenae, control of, 826; avenae, inheritance of chlamydospore characteristics in, 16; hordei, comparative studies of field collections of, 31; levis, inheritance of chlamydospore characteristics in, 16; nuda, comparative studies of field collections of, 31
Vaccinium macrocarpon Ait., transmission of false blossom through a natural graft, 670
Vascular tissue, secondary, in the bundle cap of healthy sugar-beet leaves, 694
VAUGHAN, E. K., Transmission of the crinkle disease of strawberry, 738–740
Veinbanding, in potato rugose-mosaic complex, 323, 325, 326, 330, 332, 334–337, 338; tobacco, 323–325, 326, 334
VERMILLION, M. T., see STOVER, W. G.
Verticillium, albo-atrum, pathogenicity of, 125–128; albo-atrum R. and B., pathogenicity and environal studies, 117–154; albo-atrum, susceptibility of various hosts to, 124–125; albo-atrum, temperature relations of, 121–123; "album" type, 118, 119, 121; dahliae Klebahn, pathogenicity and environal studies, 117–154; dahliae, pathogenicity of, 125–128; dahliae, susceptibility of various hosts to, 124–125; dahliae, temperature relations of, 121–123; hadromycosis, of eggplant, 130–138; hadromycosis, of potato, 143–146; hadromycosis, of tomato, 138–143; hadromycosis, pathogenicity and environal studies on, 117–154; hadromycosis, temperature relations in the field, 146–148; on elm, 21
Vigna sinensis, resistance to Rhizoctonia bataticola, 953

VINSON, CARL G., Further purification of the virus of tobacco mosaic, 35
Violet disease caused by Sphaceloma, 22–23
Virosis, complex, of tobacco, 28
Virus, acquired immunity to tobacco mosaic, 5; acquired immunity to tobacco-spot necrosis, 5; association of intracellular bodies with tobacco ring spot, 38; bean mosaic, pollen and ovule infection with, 25; cucumber mosaic, inheritability in aphids, 467–468; disease, big bud of tomato, 641–653; disease, big bud of tomato, anatomical changes in, 648; disease, big bud of tomato, transmission by grafting, 650; disease, cranberry false blossom, transmission of, 670; disease, necrotic, of potato, 32; disease, of cranberry, false blossom, 983–984; disease, peach mosaic, 17; disease, psorosis, of citrus, 930; diseases, of dahlia, 6; fig mosaic, 887; method of measuring concentration of, 752; of typical tobacco mosaic, stomatal infection with, 934–948; properties of sugar-beet mosaic, 468–470; ring-spot-like, of red clover, 746–747; seed transmission in Abutilon, 20; serologic reaction as means of determining concentration of tobacco mosaic, 4; thermal death point of sugar-beet mosaic, 470; tobacco mosaic, evidence regarding shape of particles, 34–35; tobacco mosaic, purification of, 35; tobacco mosaic, thermal death rate, 749; tobacco ring spot, increased resistance to aging in vitro from use of carbolic acid, 14–15; tolerance to dilution of sugar-beet mosaic, 470; tomato curly top, biochemical changes accompanying, 929; tomato streak, 928; tomato yellow mosaic, 34
Viruses, affecting spinach, and certain aspects of insect transmission, 446–474; causing rugose mosaic, separation of, 322–325; comparison of, on spinach, 459–466; differentiation of green and yellow mosaic, in tobacco, 10; legume mosaic, in relation to beans and peas, 562–564; of yellow mosaic, isolation of, from plants infected with tobacco mosaic, 964–974; purified plant, electrophoretic studies on, 4–5
Vitamins, effect of, on growth of fungi, 929
VOORHEES, R. K., Gibberella moniliformis on corn, 368; see EDDINS, A. H.

WADE, B. L., see ZAUMEYER, W. J.
WALKER, J. C., Comparative studies of peas resistant and susceptible to Fusarium wilt, 36; Relation of soil fertility to incidence of Aphanomyces root rot of pea, 36
WALKER, M. N., Occurrence of watermelon mosaic, 741–744
Walnut, graft disorder of, causing girdling, 408–409
WANN, F. B., see BLOOD, H. L.; and H. L. BLOOD, Biochemical changes accompanying curly top of tomato, 929
Water, bound, in corn-leaf tissues, 15; loss, in tomato mosaic, 15
WATERMAN, ALMA M., review of Insects and diseases of ornamental trees and shrubs, by E. P. FELT and W. H. RANKIN, 312–314; and MALCOLM A. MCKENZIE, A disease of Colorado fir, 108–109
Watermelon mosaic, occurrence of, 741–744
Watery soft rot on snap beans in transit, 425–428
Weather, influence of, on seedling stem blight of bean, in California, 951–953
WEBER, GEORGE F., Occurrence and pathogenicity of Nematospora spp. in Florida, 384; Pecans infected with Nematospora coryli Pegl., 1000–1001; Stem canker of Crotalaria spectabilis caused by **Diaporthe crotalariae**, n. sp., 596–604
WELCH, D. S., Nectria canker of basswood, 36
WEST, ERDMAN, Another powdery mildew on crape myrtle, 1002–1003; Powdery mildew of crape myrtle caused by **Erysiphe lagerstroemiae**, n. sp., 814–819
WESTER, R. E., see NEAL, D. C.
WESTON, WM. H., JR., A new Sclerospora from Nyasaland, 587–595; ROLAND THAXTER (1858–1932), his influence on plant pathology, 565–571
Wheat, affected by Cercosporella foot rot in Oregon, Washington, and Idaho, 33–34; bunt, average yearly losses from, in the United States, 977; bunt, losses

from, in the United States, 976; bunt, percentage of, in cars, 978; bunt, varieties resistant to, 932; Cercosporella herpotrichoides on, 33–34; disease-resistant, introduced from Russia, 8–9; effect of leaf rust on water economy and growth of, 19; flag smut, influence of soil moisture and temperature on infection by, 10–11; occurrence of Polymyxa graminis on, 20–21; pathogenicity of Wojnowicia graminis on, 1001–1002; physiologic forms of Tilletia tritici on, 16; resistance to stem rust, 25; root decoction, toxicity to Phymatotrichum omnivorum, 533–534, 536; sclerotium from, 867; stem rust, physiologic forms, in Kansas and Nebraska, 343–356; whiteheads in, ascospore discharge in Ophiobolus graminis, and its probable relation to, 721–728

WHITE, R. P., see MASSEY, L. M.

Whiteheads in wheat, ascospore discharge in Ophiobolus graminis and its probable relation to, 721–728

WHITNEY, W. A., review of Acta Phaenologica, new journal of phenology, 316–317; review of Fungous diseases of the cultivated cranberry, by C. L. SHEAR, NEIL E. STEVENS, and HENRY F. BAIN, 314–315; see LAURITZEN, J. I.

WILCOX, R. B., Relation of cranberry varieties to the spread of false blossom, 36

WILCOXON, FRANK, see MCCALLAN, S. E. A.

WILSON, E. E., Bacterial canker of Prunus spp. in California, 36–37; Development and control of olive knot, 37

WILSON, J. D., and H. A. RUNNELS, Influence of spray materials on transpiration, 37; and P. E. TILFORD, Use of formaldehyde dust with vegetable seedlings, 38

Wilt, pineapple, 207; pineapple, quick, 211; pineapple, slow, 212

Wilting of tobacco caused by Phytophthora nicotianae, 610

WINGARD, S. A., Nature of rust resistance in beans, 38

Witches' brooms on red cedar, caused by Gymnosporangium germinale, 547

Wojnowicia graminis, pathogenicity of, 1001–1002

WOLF, FREDERICK A., Roguing as a means of control of tobacco mosaic, 831–833; see BUSCH, H. J.; and E. G. MOSS, Effect of mosaic of flue-cured tobacco on yield and quality, 834–836

WOLF, FREDERICK TAYLOR, The pathology of tobacco black shank, 605–612

Wood decay as affected by temperature, 73–81

WOODROOF, NAOMI C., Two leaf spots of the peanut, 627–640

WOODS, MARK W., Intracellular bodies in ring spot, 38

WOOLMAN, HORACE MANN, biography of, 931–933

Wound overgrowth, seasonal development on apple trees in nursery, 29

WRIGHT, ERNEST, A cork-borer method for inoculating trees, 487–488

Yellows, psyllid, of tomato, 930

Yield, effect of tobacco mosaic on, 834

YOUNG, PAUL A., Sclerotinia stem canker of hollyhock, 39; Soil infestation by chlamydospores of Tilletia levis in Montana, 39

YOUNG, V. H., and C. K. MCCLELLAND, Control of oat smut, 825–830

ZAUMEYER, W. J., Transmissibility of certain legume-mosaic viruses to bean, 39; Transmission of bean-mosaic virus by insects, 40; and B. L. WADE, Mosaic diseases affecting different legumes in relation to beans and peas, 562–564

Zinc hydroxide, a substitute for calcium hydroxide in arsenical sprays, 17

Zinnia elegans, anatomy of "Kroepoek-diseased" leaf of, 175–190